

Package ‘medfate’

November 5, 2020

Type Package

Title Mediterranean Forest Simulation

Version 1.1.0

Date 2020-11-05

Description

Functions to simulate Mediterranean forest functioning and dynamics using cohort-based description of vegetation [De Cáceres et al. (2015) <doi:10.1016/j.agrformet.2015.06.012>; De Cáceres et al. (2021) <doi:10.1016/j.agrformet.2020.108233>].

License GPL (>= 2)

URL <https://vegmod.ctfc.cat/software/medfate/>

LazyLoad yes

Depends R (>= 3.5.0), sp

Imports spdep, ggplot2, methods, httr, jsonlite, meteoland (>= 0.8.1),
Rcpp (>= 0.12.12)

Suggests knitr, rmarkdown

LinkingTo Rcpp, meteoland

Encoding UTF-8

NeedsCompilation yes

VignetteBuilder utils, knitr

RoxygenNote 6.1.1

Author Miquel De Cáceres [aut, cre],
Shengli Huang [aut],
V́ctor Granda [aut],
Antoine Cabon [aut],
Jordi Mart́nez-Vilalta [ctb],
Maurizio Mencuccini [ctb],
Nicolas Martin-StPaul [ctb]

Maintainer Miquel De Cáceres <miquelcaceres@gmail.com>

Repository CRAN

Date/Publication 2020-11-05 17:30:06 UTC

R topics documented:

biophysics	3
carbon	4
conductancefunctions	6
defaultControl	8
defaultSoilParams	11
exampleforest	12
examplemeteo	13
extractSubdaily	14
fire_behaviour	15
forest	18
Forest manipulation	20
forest_mapWoodyTables	21
fuel_properties	22
growth	24
hydrology_rainInterception	28
hydrology_soilWaterInputs	29
light	31
modelInput	34
modifySpParams	39
pheno_updateLeaves	40
photo	41
Plant values	44
plot.spwb	46
plot.spwb_day	50
resetInputs	53
root	54
scalingconductance	57
SFM_metric	59
soil	61
soil hydrology	63
soil texture and hydraulics	65
soil thermodynamics	68
soilgridsParams	70
Species values	71
SpParams	72
spwb	76
spwb_day	81
spwb_ldrCalibration	84
spwb_ldrOptimization	86
spwb_maximumTranspirationRatioPlot	89
spwb_resistances	91
spwb_sensitivity	92
spwb_stress	93
spwb_validation	95
spwb_waterUseEfficiency	96
Stand values	98

supplyfunctions	99
tissuemoisture	104
transpiration	106
Vertical profiles	111

Index	113
--------------	------------

biophysics	<i>Physical and biophysical utility functions</i>
------------	---

Description

Utility functions for the calculation of biophysical variables.

Usage

```

biophysics_leafTemperature(absRad, airTemperature, u, E, leafWidth = 1.0)
biophysics_radiationDiurnalPattern(t, daylength)
biophysics_temperatureDiurnalPattern(t, tmin, tmax,
                                     tminPrev, tmaxPrev, tminNext, daylength)
biophysics_waterDynamicViscosity(temp)

```

Arguments

u	Wind speed above the leaf boundary layer (in m/s).
airTemperature	Air temperature (in °C).
tmin, tmax	Minimum and maximum daily temperature (°C).
tminPrev, tmaxPrev, tminNext	Maximum and minimum daily temperatures of the previous and following day (°C).
absRad	Absorbed long- and short-wave radiation (in W·m ⁻²).
E	Transpiration flow (in mmol H ₂ O·m ⁻² ·s ⁻¹) per one sided leaf area basis.
leafWidth	Leaf width (in cm).
t	Time of the day (in seconds).
daylength	Day length (in seconds).
temp	Temperature (°C).

Details

Function `biophysics_leafTemperature` calculates leaf temperature according to energy balance equation given in Campbell and Norman (1988). Function `biophysics_radiationDiurnalPattern` follows the equations given in Liu and Jordan (1960). Function `biophysics_temperatureDiurnalPattern` determines diurnal temperature pattern assuming a sinusoidal pattern with $T = T_{min}$ at sunrise and $T = (T_{min} + T_{max})/2$ at sunset and a linear change in temperature between sunset and T_{min} of the day after (McMurtrie et al. 1990). Function `biophysics_waterDynamicViscosity` calculates water dynamic viscosity following the Vogel (1921) equation.

Value

Values returned for each function are:

- `biophysics_leafTemperature`: leaf temperature (in °C)
- `biophysics_radiationDiurnalPattern`: the proportion of daily radiation corresponding to the input time in seconds after sunrise.
- `biophysics_temperatureDiurnalPattern`: diurnal pattern of temperature.
- `biophysics_waterDynamicViscosity`: Water dynamic viscosity relative to 20°C.

Author(s)

Miquel De Cáceres Ainsa, CTFC

References

Campbell, G. S., and J. M. Norman. 1998. An introduction to environmental biophysics: 2nd edition. (eqns. 14.1 & 14.3)

B. Y. H. Liu and R. C. Jordan, "The interrelationship and characteristic distribution of direct, diffuse and total solar radiation," *Solar Energy*, vol. 4, no. 3, pp. 1–19, 1960.

McMurtrie, R. E., D. A. Rook, and F. M. Kelliher. 1990. Modelling the yield of *Pinus radiata* on a site limited by water and nitrogen. *Forest Ecology and Management* 30:381–413.

H. Vogel, "Das Temperaturabhängigkeitsgesetz der Viskosität von Flüssigkeiten", *Physikalische Zeitschrift*, vol. 22, pp. 645–646, 1921.

See Also

[spwb](#)

carbon

Carbon-related functions

Description

Set of functions used in the calculation of carbon balance.

Usage

```
carbon_leafStarchCapacity(LAI, N, SLA, leafDensity)
carbon_leafStructuralBiomass(LAI, N, SLA)
carbon_sapwoodStarchCapacity(SA, H, L, V, woodDensity, vessel2sapwood)
carbon_sapwoodStructuralBiomass(SA, H, L, V, woodDensity)
carbon_sapwoodStructuralLivingBiomass(SA, H, L, V, woodDensity, vessel2sapwood)
carbon_sugarConcentration(osmoticWP, temp, nonSugarConc)
carbon_osmoticWaterPotential(sugarConc, temp, nonSugarConc)
carbon_relativeSapViscosity(sugarConc, temp)
carbon_sugarStarchDynamicsLeaf(sugarConc, starchConc, eqSugarConc)
```

Arguments

LAI	Leaf area index.
N	Density (ind-ha-1).
SLA	Specific leaf area (mm ² /mg = m ² /kg).
leafDensity	Density of leaf tissue (dry weight over volume).
SA	Sapwood area (cm ²).
H	Plant height (cm).
L	Coarse root length (mm) for each soil layer.
V	Proportion of fine roots in each soil layer.
woodDensity	Wood density (dry weight over volume).
vessel2sapwood	Proportion of sapwood corresponding to vessels.
osmoticWP	Osmotic water potential (MPa).
temp	Temperature (degrees Celsius).
nonSugarConc	Concentration of inorganic solutes (mol/l).
sugarConc	Concentration of soluble sugars (mol/l).
starchConc	Concentration of starch (mol/l)
eqSugarConc	Equilibrium concentration of soluble sugars (mol/l).

Value

Values returned for each function are:

- carbon_leafStarchCapacity: Capacity of storing starch in the leaf compartment (mol glucose).
- carbon_sapwoodStarchCapacity: Capacity of storing starch in the sapwood compartment (mol glucose).

Author(s)

Miquel De Cáceres Ainsa, CTFC

See Also

[growth](#)

conductancefunctions *Hydraulic conductance functions*

Description

Set of functions used in the calculation of soil and plant hydraulic conductance.

Usage

```

hydraulics_psi2K(psi, Psi_extract, ws = 3.0)
hydraulics_K2Psi(K, Psi_extract, ws = 3.0)
hydraulics_averagePsi(psi, v, c, d)
hydraulics_vulnerabilityCurvePlot(x, soil = NULL, type="leaf",
                                   psiVec = seq(-0.1, -8.0, by =-0.01),
                                   relative = FALSE, speciesNames = FALSE,
                                   draw = TRUE, ylim = NULL, xlab = NULL, ylab=NULL)
hydraulics_psiCrit(c, d, pCrit = 0.001)
hydraulics_vanGenuchtenConductance(psi, krhizomax, n, alpha)
hydraulics_xylemConductance(psi, kxylemmax, c, d)
hydraulics_xylemPsi(kxylem, kxylemmax, c, d)
hydraulics_correctConductanceForViscosity(kxylem, temp)
hydraulics_psi2Weibull(psi50, psi88)

```

Arguments

psi	A scalar (or a vector, depending on the function) with water potential (in MPa).
K	Whole-plant relative conductance (0-1).
Psi_extract	Soil water potential (in MPa) corresponding to 50% whole-plant relative conductance.
ws	Exponent of the whole-plant relative conductance Weibull function.
v	Proportion of fine roots within each soil layer.
krhizomax	Maximum rhizosphere hydraulic conductance (defined as flow per leaf surface unit and per pressure drop).
kxylemmax	Maximum xylem hydraulic conductance (defined as flow per leaf surface unit and per pressure drop).
c, d	Parameters of the Weibull function (generic xylem vulnerability curve).
n, alpha	Parameters of the Van Genuchten function (rhizosphere vulnerability curve).
kxylem	Xylem hydraulic conductance (defined as flow per surface unit and per pressure drop).
x	An object of class <code>spwbInput</code> .
soil	A list containing the description of the soil (see <code>soil</code>).
type	Plot type for <code>hydraulics_vulnerabilityCurvePlot</code> , either "leaf", "stem", "root", "rootlayer" or "rhizosphere".

psiVec	Vector of water potential values to evaluate for the vulnerability curve.
relative	A flag to relativize vulnerability curves to the [0-1] interval.
speciesNames	A flag to indicate the use of species names instead of cohort names in plots.
draw	A flag to indicate whether the vulnerability curve should be drawn or just returned.
ylim, xlab, ylab	Graphical parameters to override function defaults.
pCrit	Proportion of maximum conductance considered critical for hydraulic functioning.
psi50, psi88	Water potentials (in MPa) corresponding to 50% and 88% of percent loss of conductance.
temp	Temperature (in degrees Celsius).

Details

Details of the hydraulic model are given in a vignette. Function `hydraulics_vulnerabilityCurvePlot` draws a plot of the vulnerability curves for the given soil object and network properties of each plant cohort in `x`.

Value

Values returned for each function are:

- `hydraulics_psi2K`: Whole-plant relative conductance (0-1).
- `hydraulics_K2Psi`: Soil water potential (in MPa) corresponding to the given whole-plant relative conductance value (inverse of `hydraulics_psi2K()`).
- `hydraulics_averagePsi`: The average water potential (in MPa) across soil layers.
- `hydraulics_vanGenuchtenConductance`: Rhizosphere conductance corresponding to an input water potential (soil vulnerability curve).
- `hydraulics_xylemConductance`: Xylem conductance (flow rate per pressure drop) corresponding to an input water potential (plant vulnerability curve).
- `hydraulics_xylemPsi`: Xylem water potential (in MPa) corresponding to an input xylem conductance (flow rate per pressure drop).
- `hydraulics_psi2Weibull`: Parameters of the Weibull vulnerability curve that goes through the supplied `psi50` and `psi88` values.

Author(s)

Miquel De Cáceres Ainsa, CTFC.

References

- Sperry, J. S., F. R. Adler, G. S. Campbell, and J. P. Comstock. 1998. Limitation of plant water use by rhizosphere and xylem conductance: results from a model. *Plant, Cell & Environment* 21:347–359.
- Sperry, J. S., and D. M. Love. 2015. What plant hydraulics can tell us about responses to climate-change droughts. *New Phytologist* 207:14–27.

See Also

[hydraulics_supplyFunctionPlot](#), [hydraulics_maximumStemHydraulicConductance](#), [spwb](#), [soil](#)

Examples

```
#Manual display of vulnerability curve
kstemmax = 4 # in mmol·m-2·s-1·MPa-1
stemc = 3
stemd = -4 # in MPa
psiVec = seq(-0.1, -7.0, by =-0.01)
kstem = unlist(lapply(psiVec, hydraulics_xylemConductance, kstemmax, stemc, stemd))
plot(-psiVec, kstem, type="l",ylab="Xylem conductance (mmol·m-2·s-1·MPa-1)",
xlab="Canopy pressure (-MPa)", lwd=1.5,ylim=c(0,kstemmax))

#Load example dataset
data(exampleforestMED)
#Default species parameterization
data(SpParamsMED)

#Initialize soil with default soil params (2 layers)
examplesoil = soil(defaultSoilParams(2))

#Initialize control parameters
control = defaultControl()

#Switch to 'Sperry' transpiration mode
control$transpirationMode="Sperry"

#Initialize input
x = forest2spwbInput(exampleforestMED,examplesoil, SpParamsMED, control)

#Leaf vulnerability curves
hydraulics_vulnerabilityCurvePlot(x, type="leaf")

#Stem vulnerability curves
hydraulics_vulnerabilityCurvePlot(x, type="stem")
```

defaultControl

Default control parameters for models

Description

Creates a list with global default parameters for simulation models.

Usage

```
defaultControl()
```


Details

The function returns a list with default parameters. Users can change those defaults that need to be set to other values and use the list as input for model functions. The relevant parameters are different for each model function.

Value

A list, with the following options (default values in parentheses):

General:

- verbose (=TRUE): Boolean flag to indicate console output during calculations.
- subdailyResults (=FALSE): Boolean flag to force subdaily results to be stored (as a list called 'subdaily' of `spwb_day` objects, one by simulated date) in calls to `spwb`.
- soilFunctions (= "SX"): Soil water retention curve and conductivity functions, either 'SX' (for Saxton) or 'VG' (for Van Genuchten).
- defaultWindSpeed (= 2.5): Default wind speed value (in m/s) to be used when missing from data.

Water balance:

- snowpack (=TRUE): Boolean flag to indicate the simulation of snow accumulation and melting.
- rockyLayerDrainage (=TRUE): Boolean flag to indicate the simulation of drainage from rocky layers (> 95% of rocks).
- unlimitedSoilWater (=FALSE): Boolean flag to indicate the simulation of plant transpiration assuming that soil water is always at field capacity.
- leafPhenology (=TRUE): Boolean flag to indicate the simulation of leaf phenology for winter-deciduous species.
- unfoldingDD (=300): Degree-days for complete leaf unfolding after budburst has occurred.
- plantWaterPools (=FALSE): Boolean flag to indicate the simulation of water balance assuming that each cohort has its own water pools.
- transpirationMode (= "Granier"): Transpiration model (either 'Granier' or 'Sperry'). See `spwbInput`.
- verticalLayerSize (= 100): The size of vertical layers (in cm) for photosynthesis calculation.
- cavitationRefill (= "total"): A string indicating how refilling of embolized conduits is done ("none" - no refilling, "annual" - every first day of the year, "rate" - following a rate of new sapwood formation, "total" - instantaneous complete refilling).
- refillMaximumRate (= 0.05): Maximum rate of daily refilling of embolized conduits as sapwood area per leaf area (in $\text{cm}^2 \cdot \text{m}^{-2} \cdot \text{day}^{-1}$).

Water balance with 'Sperry':

- gainModifier, costModifier (= 1): Modifiers (exponents) of the gain and cost functions defined in Sperry et al. (2016).
- cochard (=FALSE): Boolean flag to indicate the stomatal closure when leaf water potential is below turgor loss point so that transpiration depends on cuticular properties (Martin-StPaul et al. 2017).

- ndailysteps (= 24): Number of steps into which each day is divided for determination of stomatal conductance, transpiration and photosynthesis (24 equals 1-hour intervals).
- capacitance (=FALSE): Whether the effect of plant water compartments is considered in simulations.
- klatstem (= 0.01): Stem symplastic-apoplastic lateral conductance (in $\text{mmol}\cdot\text{s}^{-1}\cdot\text{m}^{-2}\cdot\text{MPa}^{-1}$). Only relevant when capacitance = TRUE.
- klatleaf (= 0.01): Leaf symplastic-apoplastic lateral conductance (in $\text{mmol}\cdot\text{s}^{-1}\cdot\text{m}^{-2}\cdot\text{MPa}^{-1}$). Only relevant when capacitance = TRUE.
- taper (= TRUE): Whether taper of xylem conduits is accounted for when calculating aboveground stem conductance from xylem conductivity.
- numericParams: A list with the following elements:
 - * maxNsteps (= 400): Maximum number of steps in supply function.
 - * ntrial (= 200): Number of iteration trials when finding root of equation system.
 - * psiTol (= 0.0001): Tolerance value for water potential.
 - * ETol (= 0.0001): Tolerance value for flow.
- thermalCapacityLAI (=1000000): Thermal canopy capacitance per LAI unit.
- Catm (=386): Atmospheric CO₂ concentration (in $\text{micromol}\cdot\text{mol}^{-1}$ = ppm).
- fracLeafResistance (= NA): Fraction of plant total resistance (leaf+stem+root) that corresponds to leaves. This fraction is used if VCleaf_kmax = NA.
- fracRootResistance (=0.40): Fraction of plant total resistance (leaf+stem+root) that corresponds to root system.
- averageFracRhizosphereResistance (=0.15): Fraction to total continuum (leaf+stem+root+rhizosphere) resistance that corresponds to rhizosphere (averaged across soil water potential values).

Growth:

- allocationStrategy (= "Plant_kmax"): Strategy for allocation (either "Plant_kmax", for constant maximum plant conductance, or "Al2As" for constant Huber value).
- nonStomatalPhotosynthesisLimitation (=TRUE): A flag to apply limitations of photosynthesis when leaf sugar levels become too high.
- floemConductanceFactor (= 0.2): Factor to transform stem xylem conductance to stem floem conductance (only for transpirationMode = "Sperry").
- nonSugarConcentration (= 0.3): Non-sugar (inorganic) solute concentration ($\text{mol}\cdot\text{l}^{-1}$) in cells.
- equilibriumOsmoticConcentration [= c(leaf = 0.8, sapwood = 0.6)]: Equilibrium osmotic concentrations ($\text{mol}\cdot\text{l}^{-1}$) for leaf and sapwood cells. The difference helps floem transport.
- minimumSugarForGrowth [= c(leaf = 0.25, sapwood=0.25, fineroot = 0.25)]: Minimum concentration of sugars ($\text{mol}\cdot\text{l}^{-1}$) for growth to occur in different tissues.
- respirationRates [= c(leaf = 0.00260274, sapwood = 6.849315e-05, fineroot = 0.002054795)]: Maintenance respiration rates for different tissues ($\text{g gluc} \cdot \text{g dry}^{-1} \cdot \text{day}^{-1}$).
- turnoverRates [= c(sapwood = 0.0001261398, fineroot = 0.001897231)]: Turnover rates (day^{-1}) for sapwood and fineroots. Default are equivalent to 4.5% and 50% annual turnover for sapwood and fine roots, respectively.
- constructionCosts [= c(leaf = 1.5, sapwood = 1.47, fineroot = 1.30)]: Construction costs, including respiration and structural carbon, per dry weight of new tissue ($\text{g gluc} \cdot \text{g dry}^{-1}$).

- maximumRelativeGrowthRates [= c(leaf = 0.01, sapwood = 0.001, fineroot = 0.1)]: Maximum relative growth rates for leaves ($\text{m}^2 \text{ leaf} \cdot \text{cm}^{-2} \text{ sapwood} \cdot \text{day}^{-1}$), sapwood ($\text{cm}^2 \text{ sapwood} \cdot \text{cm}^{-2} \text{ sapwood} \cdot \text{day}^{-1}$) and fine roots ($\text{g dw} \cdot \text{g dw}^{-1} \cdot \text{day}^{-1}$). Values for leaves and fine roots are used for all species, whereas the value for sapwood is the default used when relative growth rate is not specified via SpParams.

Author(s)

Miquel De Cáceres Ainsa, CTFC

See Also

[spwbInput](#), [spwb](#)

defaultSoilParams	<i>Default soil parameters</i>
-------------------	--------------------------------

Description

Creates a data frame with default soil physical description for model functions.

Usage

```
defaultSoilParams(n = 4)
```

Arguments

n An integer with the number of soil layers (between two and five).

Details

The function returns a data frame with default physical soil description, with soil layers in rows. Users can change those that need to be set to other values and use the list as input for function [soil](#).

Value

A data frame with layers in rows and the following columns (and default values):

- widths (= c(300, 700, 1000, 2000)): Width of soil layers (in mm).
- clay (= 25): Clay percentage for each layer (in %).
- sand (= 25): Sand percentage for each layer (in %).
- om (= NA): Organic matter percentage for each layer (in %).
- bd (= 1.5): Bulk density for each layer (in g/cm³).
- rfc (= c(20, 40, 60, 85)): Percentage of rock fragment content (volume basis) for each layer.

Note

While this function is limited to five soil layers, user defined data frames can discretize soils using an unlimited number of soil layers.

Author(s)

Miquel De Cáceres Ainsa, CTFC

See Also

[soil](#), [soilgridsParams](#), [defaultControl](#), [SpParamsMED](#)

Examples

```
defaultSoilParams(4)
```

exampleforest

Example forest stands

Description

Data set for illustration of model behaviour. Includes a description of the plant cohorts of a forest stand.

Usage

```
data(exampleforestMED)  
data(exampleforestUS)
```

Format

An object of class [forest](#) containing the description of the tree, sapling and shrub cohorts of a forest patch.

Source

DGCN (2005). Tercer Inventario Forestal Nacional (1997-2007): Catalunya. Dirección General de Conservación de la Naturaleza, Ministerio de Medio Ambiente, Madrid.

See Also

[forest](#), [spwb](#), [forest2spwbInput](#)

Examples

```
data(exampleforestMED)
```

`examplemeteo`*Example daily meteorology data*

Description

Example data set of meteorological input.

Usage

```
data(examplemeteo)
```

Format

A data frame containing daily meteorology of a location in Catalonia (Spain) for year 2001.

MeanTemperature Mean daily temperature (in degrees Celsius).

MinTemperature Minimum daily temperature (in degrees Celsius).

MaxTemperature Maximum daily temperature (in degrees Celsius).

Precipitation Daily precipitation (in mm of water).

MeanRelativeHumidity Mean daily relative humidity (in percent).

MinRelativeHumidity Minimum daily relative humidity (in percent).

MaxRelativeHumidity Maximum daily relative humidity (in percent).

Radiation Incoming radiation (in MJ/m2).

WindSpeed Wind speed (in m/s).

WindDirection Wind direction (in degrees from North).

PET Potential evapo-transpiration (in mm of water).

Source

Interpolated from weather station data (Spanish and Catalan meteorology agencies) using package 'meteoland'.

See Also

[spwb](#)

Examples

```
data(examplemeteo)
```

extractSubdaily	<i>Extracts subdaily output</i>
-----------------	---------------------------------

Description

Given the result of simulations, this function extracts subdaily output corresponding to each simulated day and returns it as a data frame.

Usage

```
extractSubdaily(x, output = "E", dates = NULL)
```

Arguments

x	An object returned by simulation functions spwb , pwb or growth .
output	See options in section details.
dates	A date vector indicating the subset of simulated days for which subdaily output is desired.

Details

This function only works when simulations have been carried using control option 'subdailyResults = TRUE' (see [defaultControl](#)). Subdaily simulation results will then be stored as elements of the a list called 'subdaily' in the simulation output. Function `extractSubdaily` will assemble subdaily results from this list and return them as a data frame. Options for parameter 'output' are the following:

- Functions `pwb()` and `spwb()`: "E", "Ag", "An", "dEdP", "RootPsi", "StemPsi", "LeafPsi", "StemPLC", "StemRWC", "LeafRWC", "Temperature", "ExtractionInst".
- Additional options for shade and sunlit leaves in `pwb()` and `spwb()`: Either "SunlitLeaves\$x" or "ShadeLeaves\$x" where 'x' is one of the following: "Abs_SWR", "Abs_LWR", "E", "Ag", "An", "Ci", "GW", "VPD", "Tem"
- Additional options for function `growth()`: "GrossPhotosynthesis", "MaintenanceRespiration", "GrowthRespiration", "CarbonBalance", "SugarLeaf", "SugarSapwood", "StarchLeaf", "StarchSapwood", "SugarTransport".

Value

A data frame with a column 'datetime' and as many columns as plant cohorts.

Author(s)

Miquel De Cáceres Ainsa, CTFC

See Also

[spwb](#), [pwb](#), [defaultControl](#)

fire_behaviour	<i>Fire behaviour functions</i>
----------------	---------------------------------

Description

Function `fire_FCCS()` implements a modification of the fire behavior models described for the Fuel Characteristics Classification System (FCCS) in Prichard et al. (2013). Function `fire_Rothermel()` implements Rothermel's (1972) fire behaviour model (modified from package 'Rothermel' (Giorgio Vacchiano, Davide Ascoli)).

Usage

```
fire_FCCS(FCCSpropsSI, MliveSI = as.numeric(c(90, 90, 60)),
          MdeadSI = as.numeric(c(6, 6, 6, 6, 6)),
          slope = 0, windSpeedSI = 11)
fire_Rothermel(modeltype, wSI, sSI, delta, mx_dead,
               hSI, mSI, u, windDir, slope, aspect)
```

Arguments

FCCSpropsSI	A data frame describing the properties of five fuel strata (canopy, shrub, herbs, dead woody and litter) returned by <code>fuel_FCCS</code> .
MliveSI	Moisture of live fuels (in percent of dry weight) for canopy, shrub, and herb strata.
MdeadSI	Moisture of dead fuels (in percent of dry weight) for canopy, shrub, herb, woody and litter strata.
slope	Slope (in degrees).
windSpeedSI	Wind speed (in m/s) at 20 ft (6 m) over vegetation (default 11 m/s = 40 km/h)
modeltype	'S'(tatic) or 'D'(ynamic)
wSI	A vector of fuel load (t/ha) for five fuel classes.
sSI	A vector of surface-to-volume ratio (m ² /m ³) for five fuel classes.
delta	A value of fuel bed depth (cm).
mx_dead	A value of dead fuel moisture of extinction (percent).
hSI	A vector of heat content (kJ/kg) for five fuel classes.
mSI	A vector of percent moisture on a dry weight basis (percent) for five fuel classes.
u	A value of windspeed (m/s) at midflame height.
windDir	Wind direction (in degrees from north). North means blowing from north to south.
aspect	Aspect (in degrees from north).

Details

Default moisture, slope and windspeed values are benchmark conditions used to calculate fire potentials (Sandberg et al. 2007) and map vulnerability to fire.

Value

Both functions return list with fire behavior variables. In the case of `fire_FCCS`, the function returns the variables in three blocks (lists `SurfaceFire`, `CrownFire` and `FirePotentials`), and the values are:

- `SurfaceFire$`midflame_WindSpeed` [m/s]``: Midflame wind speed in the surface fire.
- `SurfaceFire$phi_wind`: Spread rate modifier due to wind.
- `SurfaceFire$phi_slope`: Spread rate modifier due to slope.
- `SurfaceFire$I_R_surf [kJ/m2/min]``: Intensity of the surface fire reaction.
- `SurfaceFire$I_R_litter [kJ/m2/min]``: Intensity of the litter fire reaction.
- `SurfaceFire$q_surf [kJ/m2]``: Heat sink of the surface fire.
- `SurfaceFire$q_litter [kJ/m2]``: Heat sink of the litter fire.
- `SurfaceFire$xi_surf`: Propagating flux ratio of the surface fire.
- `SurfaceFire$xi_litter`: Propagating flux ratio of the litter fire.
- `SurfaceFire$`ROS_surf` [m/min]``: Spread rate of the surface fire (without accounting for faster spread in the litter layer).
- `SurfaceFire$`ROS_litter` [m/min]``: Spread rate of the litter fire.
- `SurfaceFire$`ROS_windslopecap` [m/min]``: Maximum surface fire spread rate according to wind speed.
- `SurfaceFire$`ROS` [m/min]``: Final spread rate of the surface fire.
- `SurfaceFire$I_b [kW/m]``: Fireline intensity of the surface fire.
- `SurfaceFire$`FL` [m]``: Flame length of the surface fire.
- `CrownFire$I_R_canopy [kJ/m2/min]``: Intensity of the canopy fire reaction.
- `CrownFire$I_R_crown [kJ/m2/min]``: Intensity of the crown fire reaction (adding surface and canopy reactions).
- `CrownFire$q_canopy [kJ/m2]``: Heat sink of the canopy fire.
- `CrownFire$q_crown [kJ/m2]``: Heat sink of the crown fire (adding surface and canopy heat sinks).
- `CrownFire$xi_surf`: Propagating flux ratio of the crown fire.
- `CrownFire$`canopy_WindSpeed` [m/s]``: Wind speed in the canopy fire (canopy top wind speed).
- `CrownFire$WAF`: Wind speed adjustment factor for crown fires.
- `CrownFire$`ROS` [m/min]``: Spread rate of the crown fire.
- `CrownFire$Ic_ratio`: Crown initiation ratio.
- `CrownFire$I_b [kW/m]``: Fireline intensity of the crown fire.
- `CrownFire$`FL` [m]``: Flame length of the crown fire.

- FirePotentials\$RP: Surface fire reaction potential ([0-9]).
- FirePotentials\$SP: Surface fire spread rate potential ([0-9]).
- FirePotentials\$FP: Surface fire flame length potential ([0-9]).
- FirePotentials\$SFP: Surface fire potential ([0-9]).
- FirePotentials\$IC: Crown initiation potential ([0-9]).
- FirePotentials\$TC: Crown-to-crown transmission potential ([0-9]).
- FirePotentials\$RC: Crown fire spread rate potential ([0-9]).
- FirePotentials\$CFC: Crown fire potential ([0-9]).

Note

Default moisture, slope and windspeed values are benchmark conditions used to calculate fire potentials (Sandberg et al. 2007) and map vulnerability to fire.

Author(s)

Miquel De Cáceres Ainsa, CTFC

References

- Albini, F. A. (1976). Computer-based models of wildland fire behavior: A users' manual. Ogden, UT: US Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station.
- Rothermel, R. C. 1972. A mathematical model for predicting fire spread in wildland fuels. USDA Forest Service Research Paper INT USA.
- Prichard, S. J., D. V Sandberg, R. D. Ottmar, E. Eberhardt, A. Andreu, P. Eagle, and K. Swedin. 2013. Classification System Version 3.0: Technical Documentation.

See Also

[fuel_FCCS](#)

Examples

```
#Load example plot plant data
data(exampleforestMED)

#Default species parameterization
data(SpParamsMED)

#Calculate fuel properties according to FCCS
fccs = fuel_FCCS(exampleforestMED, 50,100, SpParamsMED)

#Calculate fire behavior according to FCCS
fire_FCCS(fccs)

#Load fuel model parameter data
data(SFM_metric)
```

```

#Fuel stratification (returns heights in cm)
fs = fuel_stratification(exampleforestMED, SpParamsMED)

#Correct windspeed (transform heights to m)
u = 11 #m/s
umf = u*fuel_windAdjustmentFactor(fs$surfaceLayerTopHeight/100, fs$canopyBaseHeight/100,
                                  fs$canopyTopHeight/100, 60)

#Call Rothermel function using fuel model 'A6'
fire_Rothermel(modeltype="D", wSI = as.numeric(SFM_metric["A6",2:6]),
               sSI = as.numeric(SFM_metric["A6",7:11]),
               delta = as.numeric(SFM_metric["A6",12]),
               mx_dead = as.numeric(SFM_metric["A6",13]),
               hSI = as.numeric(SFM_metric["A6",14:18]),
               mSI = c(10,10,10,30,60),
               u=umf, windDir=0, slope=0, aspect=0)

```

forest

Forest description

Description

Description of a forest stand.

Usage

```

## S3 method for class 'forest'
summary(object, SpParams, mode = "MED", detailed=FALSE, ...)
## S3 method for class 'summary.forest'
print(x, digits = getOption("digits"), ...)
emptyforest(ID="", patchsize=10000, ntree = 0, nshrub = 0)

```

Arguments

object

An object of class forest has the following structure:

- ID: An identifier of the forest stand (a string).
- patchsize: The area that forest stand description represents, in square meters.
- treeData: A data frame of tree cohorts (in rows) and the following columns:
 - Species: Non-negative integer for tree species identity (i.e., 0,1,2,...).
 - Height: Total height (in cm).
 - DBH: Diameter at breast height (in cm).
 - N: Density (number of individuals/cell).
 - Z50: Depth (in mm) corresponding to 50% of fine roots.
 - Z95: Depth (in mm) corresponding to 95% of fine roots.

- `shrubData`: A data frame of shrub cohorts (in rows) and the following columns:
 - `Species`: Non-negative integer for shrub species identity (i.e., 0,1,2,...).
 - `Height`: Total height (in cm).
 - `Cover`: Percent cover.
 - `Z50`: Depth (in mm) corresponding to 50% of fine roots.
 - `Z95`: Depth (in mm) corresponding to 95% of fine roots.
- `seedBank`: A data frame containing the abundance of seeds for each species (in rows) and the following columns:
 - `Species`: Non-negative integer for shrub species identity (i.e., 0,1,2,...).
 - `Abundance`: Abundance class (0 - none; 1 - low; 2 - medium; 3 - high; 4 - very high).
- `herbCover`: Percent cover of the herb layer.
- `herbHeight`: Mean height (in cm) of the herb layer.

<code>SpParams</code>	A data frame with species parameters (see SpParamsMED).
<code>mode</code>	Calculation mode, either "MED" or "US".
<code>detailed</code>	A boolean flag to indicate that a detailed summary is desired.
<code>x</code>	The object returned by <code>summary.forest</code> .
<code>digits</code>	Minimal number of significant digits.
<code>...</code>	Additional parameters for functions summary and print .
<code>ID</code>	An identifier of the forest stand (a string).
<code>patchsize</code>	The area of the forest stand, in square meters.
<code>ntree, nshrub</code>	Number of tree and shrub cohorts, respectively.

Details

Function `summary.forest` can be used to summarize a forest object in the console. Function `emptyforest` creates an empty forest object.

Value

Function `summary.forest` returns a list with the basal area and LAI of the forest, either expressed as totals or divided among life stages and species. Function `emptyforest` returns an empty forest object.

Author(s)

Miquel De Cáceres Ainsa, CTFC

See Also

[exampleforestMED](#), [forest_mapWoodyTables](#)

Examples

```
data(exampleforestMED)
data(SpParamsMED)

summary(exampleforestMED, SpParamsMED)
```

Forest manipulation *Forest utility functions*

Description

Functions to manipulate a [forest](#) object.

Usage

```
forest_mergeTrees(x)
```

Arguments

x An object of class [forest](#).

Value

Depending on the function:

- `forest_mergeTrees`: Another [forest](#) object with merged trees.

Author(s)

Miquel De Cáceres Ainsa, CTFC

See Also

[spwb](#), [forest](#), [summary.forest](#)

forest_mapWoodyTables *Map forest plot data*

Description

Mapping functions to facilitate building forest objects from forest plot data

Usage

```
forest_mapWoodyTables(x, y, mapping, SpParams, plot.size = NULL)
forest_mapTreeTable(x, mapping, SpParams, plot.size = NULL)
forest_mapShrubTable(y, mapping, SpParams)
```

Arguments

x	A data frame with tree records in rows and attributes in columns.
y	A data frame with shrub records in rows and attributes in columns.
mapping	A named character vector to specify mappings of columns in x or y into attributes of treeData and shrubData data frames. Accepted names (and the corresponding specifications for the columns in x and y) are: <ul style="list-style-type: none">• "Species": Species code (should follow codes in SpParams).• "Species.name": Species name. In this case, the species code will be drawn by matching names with species names in SpParams.• "N": Tree density (in ind./ha).• "plot.size": Plot size (in m2) to which each plot record refers to.• "DBH": Diameter at breast height (in cm).• "Height": Tree or shrub height (in cm).• "Z50": Depth (in mm) corresponding to 50% of fine roots.• "Z95": Depth (in mm) corresponding to 95% of fine roots.
SpParams	A data frame with species parameters (see SpParamsMED).
plot.size	The size of plot sampled area (in m2). Alternatively, 'plot.size' can a column in x and specified in mapping to indicate that trees have been measured in different subplots and, therefore, they represent different densities per hectare.

Value

Functions forest_mapTreeTable and forest_mapShrubTable return a data frame with the structure of treeData and shrubData from [forest](#) objects. Function forest_mapWoodyTable returns directly a [forest](#) object.

Author(s)

Miquel De Cáceres Ainsa, CTFC

See Also

[forest](#)

fuel_properties	<i>Fuel stratification and fuel characteristics</i>
-----------------	---

Description

Function `fuel_stratification` provides a stratification of the stand into understory and canopy strata. Function `fuel_FCCS` calculates fuel characteristics from a forest object following an adaptation of the protocols described for the Fuel Characteristics Classification System (Prichard et al. 2013). Function `fuel_windAdjustmentFactor` determines the adjustment factor of wind for surface fires, according to Andrews (2012).

Usage

```
fuel_stratification(object, SpParams, gdd = NA, mode = "MED",
                   heightProfileStep = 10.0, maxHeightProfile = 5000.0,
                   bulkDensityThreshold = 0.05)
fuel_FCCS(object, ShrubCover, CanopyCover, SpParams, cohortFMC = as.numeric(c()),
           gdd = NA, mode = "MED", heightProfileStep = 10, maxHeightProfile = 5000,
           bulkDensityThreshold = 0.05)
fuel_windAdjustmentFactor(topShrubHeight, bottomCanopyHeight, topCanopyHeight,
                          canopyCover)
```

Arguments

<code>object</code>	An object of class <code>forest</code>
<code>ShrubCover</code>	Total shrub cover (in percent) of the stand.
<code>CanopyCover</code>	Total canopy cover (in percent) of the stand.
<code>SpParams</code>	A data frame with species parameters (see <code>SpParamsMED</code>).
<code>cohortFMC</code>	A numeric vector of (actual) fuel moisture content by cohort (e.g. taken from the result of <code>fuel_cohortFineFMC</code>).
<code>gdd</code>	Growth degree-days.
<code>mode</code>	Calculation mode, either "MED" or "US".
<code>heightProfileStep</code>	Precision for the fuel bulk density profile.
<code>maxHeightProfile</code>	Maximum height for the fuel bulk density profile.
<code>bulkDensityThreshold</code>	Minimum fuel bulk density to delimit fuel strata.
<code>topShrubHeight</code>	Shrub stratum top height (in m).
<code>bottomCanopyHeight</code>	Canopy base height (in m).
<code>topCanopyHeight</code>	Canopy top height (in m).
<code>canopyCover</code>	Canopy percent cover.

Details

Details are described in a vignette.

Value

Function `fuel_FCCS` returns a data frame with five rows corresponding to fuel layers: canopy, shrub, herb, woody and litter. Columns correspond fuel properties:

- `w`: Fine fuel loading (in kg/m²).
- `cover`: Percent cover.
- `hbc`: Height to base of crowns (in m).
- `htc`: Height to top of crowns (in m).
- `delta`: Fuel depth (in m).
- `rhob`: Fuel bulk density (in kg/m³).
- `rhop`: Fuel particle density (in kg/m³).
- `PV`: Particle volume (in m³/m²).
- `beta`: Packing ratio (unitless).
- `betarel`: Relative packing ratio (unitless).
- `etabetarel`: Reaction efficiency (unitless).
- `sigma`: Surface area-to-volume ratio (m²/m³).
- `pDead`: Proportion of dead fuels.
- `FAI`: Fuel area index (unitless).
- `h`: High heat content (in kJ/kg).
- `etaF`: Flammability modifier (between 1 and 2).
- `RV`: Reactive volume (in m³/m²).
- `MinFMC`: Minimum fuel moisture content (as percent over dry weight).
- `MaxFMC`: Maximum fuel moisture content (as percent over dry weight).

Function `fuel_stratification` returns a list with the following items:

- `surfaceLayerBaseHeight`: Base height of crowns of shrubs in the surface layer (in cm).
- `surfaceLayerTopHeight`: Top height of crowns of shrubs in the surface layer (in cm).
- `understoryLAI`: Cumulated LAI of the understory layer (i.e. leaf area comprised between surface layer base and top heights).
- `canopyBaseHeight`: Base height of tree crowns in the canopy (in cm).
- `canopyTopHeight`: Top height of tree crowns in the canopy (in cm).
- `canopyLAI`: Cumulated LAI of the canopy (i.e. leaf area comprised between canopy base and top heights).

Function `fuel_cohortFineFMC` returns a list with three matrices (for leaves, twigs and fine fuels). Each of them contains live moisture content values for each day (in rows) and plant cohort (in columns).

Function `fuel_windAdjustmentFactor` returns a value between 0 and 1.

Author(s)

Miquel De Cáceres Ainsa, CTFC

References

Andrews, P. L. 2012. Modeling wind adjustment factor and midflame wind speed for Rothermel's surface fire spread model. USDA Forest Service - General Technical Report RMRS-GTR:1-39.

Prichard, S. J., D. V Sandberg, R. D. Ottmar, E. Eberhardt, A. Andreu, P. Eagle, and K. Swedin. 2013. Classification System Version 3.0: Technical Documentation.

Reinhardt, E., D. Lutes, and J. Scott. 2006. FuelCalc: A method for estimating fuel characteristics. Pages 273-282.

See Also

[fire_FCCS](#), [spwb](#)

Examples

```
#Load example plot plant data
data(exampleforestMED)

#Default species parameterization
data(SpParamsMED)

#Show stratification of fuels
fuel_stratification(exampleforestMED, SpParamsMED)

#Calculate fuel properties according to FCCS
fccs = fuel_FCCS(exampleforestMED, 50,100, SpParamsMED)
fccs

fuel_windAdjustmentFactor(fccs$htc[2], fccs$hbc[1], fccs$htc[1], fccs$cover[1])
```

growth

Forest growth

Description

Function growth is a forest growth model that calculates changes in leaf area, sapwood area and structural variables for all plant cohorts in a given forest stand during a period specified in the input climatic data.

Usage

```
growth(x, soil, meteo, latitude, elevation = NA, slope = NA, aspect = NA)
```


Arguments

x	An object of class <code>growthInput</code> .
soil	A list containing the description of the soil (see <code>soil</code>).
meteo	A data frame with daily meteorological data series. Row names of the data frame should correspond to date strings with format "yyyy-mm-dd" (see <code>Date</code>). When <code>x\$TranspirationMode = "Granier"</code> the following columns are required: <ul style="list-style-type: none"> • <code>MeanTemperature</code>: Mean temperature (in degrees Celsius). • <code>Precipitation</code>: Precipitation (in mm). • <code>Radiation</code>: Solar radiation (in MJ/m²/day), required only if <code>snowpack = TRUE</code>. • <code>PET</code>: Potential evapotranspiration (in mm). • <code>WindSpeed</code>: Wind speed (in m/s). If not available, this column can be left with NA values. <p>When <code>x\$TranspirationMode = "Sperry"</code> the following columns are required:</p> <ul style="list-style-type: none"> • <code>MeanTemperature</code>: Mean temperature (in degrees Celsius). • <code>MinTemperature</code>: Minimum temperature (in degrees Celsius). • <code>MaxTemperature</code>: Maximum temperature (in degrees Celsius). • <code>MinRelativeHumidity</code>: Minimum relative humidity (in percent). • <code>MaxRelativeHumidity</code>: Maximum relative humidity (in percent). • <code>Precipitation</code>: Precipitation (in mm). • <code>Radiation</code>: Solar radiation (in MJ/m²/day). • <code>WindSpeed</code>: Wind speed (in m/s). If not available, this column can be left with NA values.
latitude	Latitude (in degrees). Required when <code>x\$TranspirationMode = "Sperry"</code> .
elevation, slope, aspect	Elevation above sea level (in m), slope (in degrees) and aspect (in degrees from North). Required when <code>x\$TranspirationMode = "Sperry"</code> . Elevation is also required for 'Granier' if snowpack dynamics are simulated.

Details

Detailed model description is available in the vignettes section. Simulations using the 'Sperry' transpiration mode are computationally much more expensive than those using the simple transpiration mode.

Value

A list of class 'growth' with the following elements:

- `"latitude"`: Latitude (in degrees) given as input.
- `"topography"`: Vector with elevation, slope and aspect given as input.
- `"growthInput"`: A copy of the object x of class `growthInput` given as input.
- `"soilInput"`: A copy of the object soil of class `soil` given as input.
- `"WaterBalance"`: A data frame where different water balance variables (see `spwb`).

- "EnergyBalance": A data frame with the daily values of energy balance components for the soil and the canopy (only for `transpirationMode = "Sperry"`; see [spwb](#)).
- "Temperature": A data frame with the daily values of minimum/mean/maximum temperatures for the atmosphere (input), canopy and soil (only for `transpirationMode = "Sperry"`; see [spwb](#)).
- "Soil": A data frame where different soil variables (see [spwb](#)).
- "Stand": A data frame where different stand-level variables (see [spwb](#)).
- "Plants": A list of daily results for plant cohorts (see [spwb](#)).
- "SunlitLeaves" and "ShadeLeaves": A list with daily results for sunlit and shade leaves (only for `transpirationMode = "Sperry"`; see [spwb](#)).
- "PlantCarbonBalance": A list of daily carbon balance results for plant cohorts, with elements:
 - "GrossPhotosynthesis": Gross photosynthesis rate per dry weight of living biomass ($\text{g gluc} \cdot \text{g dry-1}$).
 - "MaintenanceRespiration": Maintenance respiration rate per dry weight of living biomass ($\text{g gluc} \cdot \text{g dry-1}$).
 - "GrowthRespiration": Growth respiration rate per dry weight of living biomass ($\text{g gluc} \cdot \text{g dry-1}$).
 - "CarbonBalance": Carbon balance (photosynthesis - respiration) per dry weight of living biomass ($\text{g gluc} \cdot \text{g dry-1}$).
 - "SugarLeaf": Sugar concentration ($\text{mol}\cdot\text{l-1}$) in leaves.
 - "StarchLeaf": Starch concentration ($\text{mol}\cdot\text{l-1}$) in leaves.
 - "SugarSapwood": Sugar concentration ($\text{mol}\cdot\text{l-1}$) in sapwood.
 - "StarchSapwood": Starch concentration ($\text{mol}\cdot\text{l-1}$) in sapwood.
- "PlantGrowth": A list of daily growth results for plant cohorts, with elements:
 - "SapwoodArea": Daily amount of sapwood area (in cm^2) for an average individual of each plant cohort.
 - "LeafArea": Daily amount of leaf area (in m^2) for an average individual of each plant cohort.
 - "FineRootArea": Daily amount of fine root area (in m^2) for an average individual of each plant cohort (only for `transpirationMode = "Sperry"`).
 - "SAGrowth": Sapwood area relative growth rate (in $\text{cm}^2\cdot\text{cm-2}\cdot\text{day-1}$) for an average individual of each plant cohort.
 - "LAGrowth": Leaf area growth relative to sapwood area (in $\text{m}^2\cdot\text{cm-2}\cdot\text{day-1}$) for an average individual of each plant cohort.
 - "FRAGrowth": Fine root area growth relative to sapwood area (in $\text{m}^2\cdot\text{cm-2}\cdot\text{day-1}$) for an average individual of each plant cohort (only for `transpirationMode = "Sperry"`).
- "StandStructures": A list with the aboveground structure of the stand at the beginning of the simulation and after each simulated year.
- "StandSummary": A data frame with stand-level summaries (leaf area index, tree basal area, tree density, shrub cover, etc.) at the beginning of the simulation and after each simulated year.
- "subdaily": A list of objects of class [growth_day](#), one per day simulated (only if required in control parameters, see [defaultControl](#)).

Note

Objects `x` and `soil` are modified during the simulation.

Author(s)

Miquel De Cáceres Ainsa, CTFC

See Also

[growthInput](#), [growth_day](#), [plot.growth](#)

Examples

```
#Load example daily meteorological data
data(examplemeteo)

#Load example plot plant data
data(exampleforestMED)

#Default species parameterization
data(SpParamsMED)

#Initialize control parameters
control = defaultControl()

#Initialize soil with default soil params (4 layers)
examplesoil1 = soil(defaultSoilParams(4))

#Initialize vegetation input
x1 = forest2growthInput(exampleforestMED, examplesoil1, SpParamsMED, control)

#Call simulation function
G1<-growth(x1, examplesoil1, examplemeteo, latitude = 41.82592, elevation = 100)

## Not run:
#Switch to 'Sperry' transpiration mode
control$transpirationMode="Sperry"

#Initialize soil with default soil params (4 layers)
examplesoil2 = soil(defaultSoilParams(4))

#Initialize vegetation input
x2 = forest2growthInput(exampleforestMED,examplesoil2, SpParamsMED, control)

#Call simulation function (11 days)
d = 100:110
G2<-growth(x2, examplesoil2, examplemeteo[d,], latitude = 41.82592, elevation = 100)

## End(Not run)
```

hydrology_rainInterception

Rainfall interception

Description

Function `hydrology_rainInterception` calculates the amount of rainfall intercepted daily by the canopy, given a rainfall and canopy characteristics. Two canopy interception models are currently available: the sparse Gash (1995) model and the Liu (2001) model. In both cases the current implementation assumes no trunk interception.

Usage

```
hydrology_rainInterception(Rainfall, Cm, p, ER=0.05, method="Gash1995")
hydrology_erFactor(doy, pet, prec, Rconv = 5.6, Rsyn = 1.5)
hydrology_interceptionPlot(x, SpParams, ER = 0.05, gdd = NA, throughfall = FALSE)
```

Arguments

Rainfall	A numeric vector of (daily) rainfall.
Cm	Canopy water storage capacity.
p	Proportion of throughfall (normally $1 - c$, where c is the canopy cover).
ER	The ratio of evaporation rate to rainfall rate.
method	Rainfall interception method (either "Gash1995" or "Liu2001").
doy	Day of the year.
pet	Potential evapotranspiration for a given day (mm).
prec	Precipitation for a given day (mm).
Rconv, Rsyn	Rainfall rate for convective storms and synoptic storms, respectively, in mm/h.
x	An object of class <code>spwbInput</code> .
SpParams	A data frame with species parameters (see <code>SpParamsMED</code> and <code>SpParamsMED</code>).
gdd	Growth degree days (in Celsius).
throughfall	Boolean flag to plot relative throughfall instead of percentage of intercepted rainfall.

Details

Function `hydrology_rainInterception` can accept either vectors or scalars as parameters `Cm`, `p` and `ER`. If they are supplied as vectors they should be of the same length as `Rainfall`.

Function `hydrology_erFactor` calculates the evaporation-to-rainfall ratio for input values of potential evapotranspiration and rainfall, while accounting for seasonal variation in rainfall intensity (mm/h). Default values `Rconv = 5.6` and `Rsyn = 1.5` come from Miralles et al. (2010).

Value

Function hydrology_rainInterception returns a vector of the same length as Rainfall containing intercepted rain values. Function hydrology_erFactor returns a scalar with the evaporation-to-rainfall ratio.

Author(s)

Miquel De Cáceres Ainsa, CTFC

References

Liu (2001). Evaluation of the Liu model for predicting rainfall interception in forests world-wide. - Hydrol. Process. 15: 2341-2360.

Gash (1979). An analytical model of rainfall interception by forests. - Quarterly Journal of the Royal Meteorological Society.

Gash et al. (1995). Estimating sparse forest rainfall interception with an analytical model. - Journal of Hydrology.

Miralles DG, Gash JH, Holmes TRH, et al (2010) Global canopy interception from satellite observations. J Geophys Res 115:D16122. doi: 10.1029/2009JD013530.

See Also

[spwb](#)

Examples

```
#Load example plot plant data
data(exampleforestMED)

#Default species parameterization
data(SpParamsMED)

#Draw rainfall interception for two values of the E/R ratio
hydrology_interceptionPlot(exampleforestMED, SpParamsMED, ER = c(0.05, 0.2))
```

hydrology_soilWaterInputs

Soil water processes

Description

High-level functions for hydrological processes. Function hydrology_soilWaterInputs performs canopy water interception and snow accumulation/melt. Function hydrology_soilInfiltrationPercolation performs soil infiltration and percolation from the input given by the previous function.

Usage

```
hydrology_soilWaterInputs(soil, soilFunctions, prec, er, tday, rad,
                          elevation, Cm, LgroundPAR, LgroundSWR, runon = 0,
                          snowpack = TRUE, modifySoil = TRUE)
hydrology_soilInfiltrationPercolation(soil, soilFunctions, waterInput,
                                      rockyLayerDrainage = TRUE, modifySoil = TRUE)
```

Arguments

soil	A list containing the description of the soil (see soil).
soilFunctions	Soil water retention curve and conductivity functions, either 'SX' (for Saxton) or 'VG' (for Van Genuchten).
prec	Precipitation for a given day (mm)
waterInput	Soil water input for a given day (mm).
er	The ratio of evaporation rate to rainfall rate.
tday	Average day temperature (°C).
rad	Solar radiation (in MJ/m ² /day).
elevation	Altitude above sea level (m).
Cm	Canopy water storage capacity.
LgroundPAR	Proportion of photosynthetically-active radiation (PAR) reaching the ground.
LgroundSWR	Proportion of short-wave radiation (SWR) reaching the ground.
runon	Surface water amount running on the target area from upslope (in mm).
snowpack	Boolean flag to indicate the simulation of snow accumulation and melting.
rockyLayerDrainage	Boolean flag to indicate the simulation of drainage from rocky layers (> 95% of rocks).
modifySoil	Boolean flag to indicate that the input soil object should be modified during the simulation.

Details

The function simulates different vertical hydrological processes, which are described separately in other functions. If `modifySoil = TRUE` the function will modify the soil object (including both soil moisture and the snowpack on its surface) as a result of simulating hydrological processes.

Value

Function `hydrology_soilWaterInputs` returns a named vector with the following elements, all in mm:

Rain	Precipitation as rainfall.
Snow	Precipitation as snow.
Interception	Rainfall water intercepted by the canopy and evaporated.
NetRain	Rainfall reaching the ground.

Snowmelt	Snow melted during the day, and added to the water infiltrated.
Runon	Surface water amount running on the target area from upslope.
Input	Total soil input, including runon, snowmelt and net rain.

Function `hydrology_soilInfiltrationPercolation` returns a named vector with the following elements, all in mm:

Infiltration	Water infiltrated into the soil (i.e. throughfall + runon + snowmelt - runoff).
Runoff	Surface water leaving the target area.
DeepDrainage	Water leaving the target soil towards the water table.

Author(s)

Miquel De Cáceres Ainsa, CTFC

See Also

[spwb_day](#), [hydrology_rainInterception](#), [hydrology_soilEvaporation](#)

light

Light extinction and absorption functions

Description

Functions `light_layerIrradianceFraction` and `light_layerIrradianceFractionBottomUp` calculate the fraction of above-canopy irradiance (and the soil irradiance, respectively) reaching each vegetation layer. Function `light_layerSunlitFraction` calculates the proportion of sunlit leaves in each vegetation layer. Function `light_cohortSunlitShadeAbsorbedRadiation` calculates the amount of radiation absorbed by cohort and vegetation layers, while differentiating between sunlit and shade leaves.

Usage

```
light_layerIrradianceFraction(LAIme, LAImd, LAImx, k, alpha,
                             trunkExtinctionFraction = 0.1)
light_layerIrradianceFractionBottomUp(LAIme, LAImd, LAImx, k, alpha,
                                       trunkExtinctionFraction = 0.1)
light_layerSunlitFraction(LAIme, LAImd, kb)
light_cohortSunlitShadeAbsorbedRadiation(Ib0, Id0, Ibf, Idf, beta,
                                         LAIme, LAImd,
                                         kb, kd, alpha, gamma)
light_instantaneousLightExtinctionAbsortion(LAIme, LAImd, LAImx,
                                             kPAR, alphaSWR, gammaSWR,
                                             ddd, LWR_diffuse,
                                             nimesteps = 24, canopyMode = "sunshade",
                                             trunkExtinctionFraction = 0.1)
light_cohortAbsorbedSWRFraction(z, x, SpParams, gdd = NA)
```

Arguments

LAI _{live}	A numeric matrix of live expanded LAI values per vegetation layer (row) and cohort (column).
LAI _{dead}	A numeric matrix of dead LAI values per vegetation layer (row) and cohort (column).
LAI _{max}	A numeric matrix of maximum LAI values per vegetation layer (row) and cohort (column).
k	A vector of light extinction coefficients.
k _b	A vector of direct light extinction coefficients.
k _d	A vector of diffuse light extinction coefficients.
I _{b0}	Above-canopy direct incident radiation.
I _{d0}	Above-canopy diffuse incident radiation.
I _b _f	Fraction of above-canopy direct radiation reaching each vegetation layer.
I _d _f	Fraction of above-canopy diffuse radiation reaching each vegetation layer.
alpha	A vector of leaf absorbance by species.
beta	Solar elevation (in radians).
gamma	Vector of canopy reflectance values.
k _{PAR}	A vector of visible light extinction coefficients for each cohort.
alpha _{SWR}	A vector of short-wave absorbance coefficients for each cohort.
gamma _{SWR}	A vector of short-wave reflectance coefficients (albedo) for each cohort.
ddd	A dataframe with direct and diffuse radiation for different subdaily time steps (see function <code>radiation_directDiffuseDay</code> in package <code>meteoland</code>).
LWR _{diffuse}	A vector with diffuse longwave radiation for different subdaily time steps.
ntimesteps	Number of subdaily time steps.
canopyMode	Indicates how crowns should be described to calculate photosynthesis/extinction. Accepted values are "sunshade" (distinguishes photosynthesis in sun leaves from shade leaves) and "multilayer" (distinguishes photosynthesis of sun leaves and shade leaves in each canopy layer).
trunkExtinctionFraction	Fraction of extinction due to trunks (for winter deciduous forests).
x	An object of class <code>forest</code>
SpParams	A data frame with species parameters (see <code>SpParamsMED</code>).
z	A numeric vector with height values.
gdd	Growth degree days.

Details

Function codification adapted from Anten & Bastiaans (2016). Vegetation layers are assumed to be ordered from bottom to top.

Value

Functions `light_layerIrradianceFraction`, `light_layerIrradianceFractionBottomUp` and `light_layerSunlitFraction` return a numeric vector of length equal to the number of vegetation layers. Function `light_cohortSunlitShadeAbsorbedRadiation` returns a list with two elements (matrices): `I_sunlit` and `I_shade`.

Author(s)

Miquel De Cáceres Ainsa, CTFC

References

Anten, N.P.R., Bastiaans, L., 2016. The use of canopy models to analyze light competition among plants, in: Hikosaka, K., Niinemets, U., Anten, N.P.R. (Eds.), *Canopy Photosynthesis: From Basics to Application*. Springer, pp. 379–398.

See Also

[spwb](#)

Examples

```
LAI = 2
nlayer = 10
LAIlayerlive = matrix(rep(LAI/nlayer,nlayer),nlayer,1)
LAIlayerdead = matrix(0,nlayer,1)
kb = 0.8
kd_PAR = 0.5
kd_SWR = kd_PAR/1.35
alpha_PAR = 0.9
gamma_PAR = 0.04
gamma_SWR = 0.05
alpha_SWR = 0.7

Ibfp = light_layerIrradianceFraction(LAIlayerlive,LAIlayerdead,LAIlayerlive,kb, alpha_PAR)
Idfp = light_layerIrradianceFraction(LAIlayerlive,LAIlayerdead,LAIlayerlive,kd_PAR, alpha_PAR)
Ibfsr = light_layerIrradianceFraction(LAIlayerlive,LAIlayerdead,LAIlayerlive,kb, alpha_SWR)
Idfsr = light_layerIrradianceFraction(LAIlayerlive,LAIlayerdead,LAIlayerlive,kd_SWR, alpha_SWR)
fsunlit = light_layerSunlitFraction(LAIlayerlive, LAIlayerdead, kb)
SHarea = (1-fsunlit)*LAIlayerlive[,1]
SLarea = fsunlit*LAIlayerlive[,1]

par(mar=c(4,4,1,1), mfrow=c(1,2))
plot(Ibfp*100, 1:nlayer,type="l", ylab="Layer",
      xlab="Percentage of irradiance", xlim=c(0,100), ylim=c(1,nlayer), col="dark green")
lines(Idfp*100, 1:nlayer, col="dark green", lty=2)
lines(Ibfsr*100, 1:nlayer, col="red")
lines(Idfsr*100, 1:nlayer, col="red", lty=2)

plot(fsunlit*100, 1:nlayer,type="l", ylab="Layer",
      xlab="Percentage of leaves", xlim=c(0,100), ylim=c(1,nlayer))
```

```

lines((1-fsunlit)*100, 1:nlayer, lty=2)

solarElevation = 0.67
SWR_direct = 1100
SWR_diffuse = 300
PAR_direct = 550
PAR_diffuse = 150

abs_PAR = light_cohortSunlitShadeAbsorbedRadiation(PAR_direct, PAR_diffuse,
            Ibfpar, Idfpar, beta = solarElevation,
            LAIlayerlive, LAIlayerdead, kb, kd_PAR, alpha_PAR, gamma_PAR)
abs_SWR = light_cohortSunlitShadeAbsorbedRadiation(SWR_direct, SWR_diffuse,
            Ibfswr, Idfswr, beta = solarElevation,
            LAIlayerlive, LAIlayerdead, kb, kd_SWR, alpha_SWR, gamma_SWR)
par(mar=c(4,4,1,1), mfrow=c(1,2))
absRadSL = abs_SWR$I_sunlit[,1]
absRadSH = abs_SWR$I_shade[,1]
lambda = 546.6507
QSL = abs_PAR$I_sunlit[,1]*lambda*0.836*0.01
QSH = abs_PAR$I_shade[,1]*lambda*0.836*0.01
plot(QSL, 1:nlayer,type="l", ylab="Layer",
      xlab="Absorbed PAR quantum flux per leaf area", ylim=c(1,nlayer), col="dark green",
      xlim=c(0,max(QSL)))
lines(QSH, 1:nlayer, col="dark green", lty=2)
plot(absRadSL, 1:nlayer,type="l", ylab="Layer",
      xlab="Absorbed SWR per leaf area (W/m2)", ylim=c(1,nlayer), col="red",
      xlim=c(0,max(absRadSL)))
lines(absRadSH, 1:nlayer, col="red", lty=2)

```

modelInput

Input for simulation models

Description

Functions `forest2spwbInput` and `forest2growthInput` take an object of class `forest` and calculate input data for functions `spwb`, `pwb` and `growth`, respectively. Functions `spwbInput` and `growthInput` do the same but starting from different input data. Function `forest2aboveground` calculates aboveground variables that may be used in `spwbInput` and `growthInput` functions. Function `forest2belowground` calculates belowground fine root distribution.

Usage

```

forest2aboveground(x, SpParams, gdd = NA, mode = "MED")
forest2belowground(x, soil)
forest2growthInput(x, soil, SpParams, control)
forest2spwbInput(x, soil, SpParams, control, mode = "MED")
growthInput(above, Z50, Z95, soil, SpParams, control)
spwbInput(above, Z50, Z95, soil, SpParams, control)

```

Arguments

x	An object of class <code>forest</code> .
SpParams	A data frame with species parameters (see <code>SpParamsMED</code> and <code>SpParamsMED</code>).
gdd	Growth degree days to account for leaf phenology effects (in Celsius). This should be left NA in most applications.
mode	Calculation mode, either "MED" or "US".
soil	An object of class <code>soil</code> .
control	A list with default control parameters (see <code>defaultControl</code>).
above	A data frame with aboveground plant information (see the return value of <code>forest2aboveground</code> below). In the case of <code>spwbInput</code> the variables should include SP, N, LAI_live, LAI_dead, H and CR. In the case of <code>growthInput</code> variables should include DBH and Cover.
Z50, Z95	Numeric vectors with cohort depths (in mm) corresponding to 50% and 95% of fine roots.

Details

Functions `forest2spwbInput` and `forest2abovegroundInput` extracts height and species identity from plant cohorts of `x`, and calculate leaf area index and crown ratio. `forest2spwbInput` also calculates the distribution of fine roots across soil. Both `forest2spwbInput` and `spwbInput` find parameter values for each plant cohort according to the parameters of its species as specified in `SpParams`. If `control$transpirationMode = "Sperry"` the functions also estimate the maximum conductance of rhizosphere, root xylem and stem xylem elements.

Value

Function `forest2aboveground()` returns a data frame with the following columns (rows are identified as specified by function `plant_ID`):

- SP: Species identity (an integer) (first species is 0).
- N: Cohort density (ind/ha) (see function `plant_density`).
- DBH: Tree diameter at breast height (cm).
- H: Plant total height (cm).
- CR: Crown ratio (crown length to total height) (between 0 and 1).
- LAI_live: Live leaf area index (m²/m²) (one-side leaf area relative to plot area), includes leaves in winter dormant buds.
- LAI_expanded: Leaf area index of expanded leaves (m²/m²) (one-side leaf area relative to plot area).
- LAI_dead: Dead leaf area index (m²/m²) (one-side leaf area relative to plot area).

Functions `forest2spwbInput()` and `spwbInput()` return a list of class `spwbInput` with the following elements (rows of data frames are identified as specified by function `plant_ID`):

- `control`: List with control parameters (see `defaultControl`).
- `canopy`: A list of stand-level state variables.

- cohorts: A data frame with cohort information, with columns SP and Name.
 - above: A data frame with columns H, CR and LAI (see function forest2aboveground).
 - below: A data frame with columns Z50, Z95. If control\$transpirationMode = "Sperry" additional columns are fineRootBiomass and coarseRootSoilVolume.
 - belowLayers: A list. If control\$transpirationMode = "Granier" it contains elements:
 - V: A matrix with the proportion of fine roots of each cohort (in rows) in each soil layer (in columns).
 - L: A matrix with the length of coarse roots of each cohort (in rows) in each soil layer (in columns).
 - Wpool: A matrix with the soil moisture relative to field capacity around the rhizosphere of each cohort (in rows) in each soil layer (in columns).
- If control\$transpirationMode = "Sperry" there are the following additional elements:
- VGrhizo_kmax: A matrix with maximum rhizosphere conductance values of each cohort (in rows) in each soil layer (in columns).
 - VGroot_kmax: A matrix with maximum root xylem conductance values of each cohort (in rows) in each soil layer (in columns).
 - RhizoPsi: A matrix with the water potential around the rhizosphere of each cohort (in rows) in each soil layer (in columns).
- paramsPhenology: A data frame with leaf phenology parameters:
 - Sgdd: Growth degree days needed for leaf budburst (for winter deciduous species).
 - paramsAnatomy: A data frame with plant anatomy parameters for each cohort (only if control\$transpirationMode = "Sperry"):
 - Hmax: Maximum plant height (cm).
 - Hmed: Median plant height (cm).
 - A12As: Leaf area to sapwood area ratio (in m²·m⁻²).
 - SLA: Specific leaf area (mm²/mg = m²/kg).
 - LeafWidth: Leaf width (in cm).
 - LeafDensity: Density of leaf tissue (dry weight over volume).
 - WoodDensity: Density of wood tissue (dry weight over volume).
 - FineRootDensity: Density of fine root tissue (dry weight over volume).
 - SRL: Specific Root length (cm·g⁻¹).
 - RLD: Root length density (cm·cm⁻³).
 - r635: Ratio between the weight of leaves plus branches and the weight of leaves alone for branches of 6.35 mm.
 - paramsInterception: A data frame with rain interception and light extinction parameters for each cohort:
 - kPAR: PAR extinction coefficient.
 - g: Canopy water retention capacity per LAI unit (mm/LAI).
- If control\$transpirationMode = "Sperry" additional columns are:
- gammaSWR: Reflectance (albedo) coefficient for SWR .
 - alphaSWR: Absorbance coefficient for SWR .

- `paramsTranspiration`: A data frame with parameters for transpiration and photosynthesis. If `control$transpirationMode = "Granier"`, columns are:
 - `Psi_Extract`: Water potential corresponding to 50% relative conductance (in MPa).
 - `WUE`: Water use efficiency for carbon assimilation (g C/mm water).
 - `pRootDisc`: Proportion of whole-plant conductance leading to disconnection from soil.
 If `control$transpirationMode = "Sperry"` columns are:
 - `Gwmax`: Maximum stomatal conductance to water vapor (in mol H₂O·m⁻²·s⁻¹).
 - `Vmax298`: Maximum Rubisco carboxylation rate at 25°C (in micromol CO₂·s⁻¹·m⁻²).
 - `Jmax298`: Maximum rate of electron transport at 25°C (in micromol photons·s⁻¹·m⁻²).
 - `Kmax_stemxylem`: Sapwood-specific hydraulic conductivity of stem xylem (in kg H₂O·s⁻¹·m⁻²).
 - `Kmax_rootxylem`: Sapwood-specific hydraulic conductivity of root xylem (in kg H₂O·s⁻¹·m⁻²).
 - `VCleaf_kmax`: Maximum leaf hydraulic conductance.
 - `VCleaf_c`, `VCleaf_d`: Parameters of the leaf vulnerability curve.
 - `VCstem_kmax`: Maximum stem xylem conductance.
 - `VCstem_c`, `VCstem_d`: Parameters of the stem xylem vulnerability curve.
 - `VCroot_c`, `VCroot_d`: Parameters of the root xylem vulnerability curve.
 - `Plant_kmax`: Maximum whole-plant conductance.
- `paramsWaterStorage`: A data frame with plant water storage parameters for each cohort (only if `control$transpirationMode = "Sperry"`):
 - `LeafPI0`: Osmotic potential at full turgor of leaves (MPa).
 - `LeafEPS`: Modulus of elasticity (capacity of the cell wall to resist changes in volume in response to changes in turgor) of leaves (MPa).
 - `LeafAF`: Apoplastic fraction (proportion of water outside the living cells) in leaves.
 - `Vleaf`: Storage water capacity in leaves, per leaf area (L/m²).
 - `StemPI0`: Osmotic potential at full turgor of symplastic xylem tissue (MPa).
 - `StemEPS`: Modulus of elasticity (capacity of the cell wall to resist changes in volume in response to changes in turgor) of symplastic xylem tissue (Mpa).
 - `StemAF`: Apoplastic fraction (proportion of water outside the living cells) in stem xylem.
 - `Vstem`: Storage water capacity in sapwood, per leaf area (L/m²).
- `internalPhenology` and `internalWater`: data frames to store internal state variables.

Functions `forest2growthInput` and `growthInput` return a list of class `growthInput` with the same elements as `spwbInput`, but with additional information.

- Element above includes the following additional columns:
 - `LA_live`: Live leaf area per individual (m²/ind).
 - `LA_dead`: Dead leaf area per individual (m²/ind).
 - `SA`: Live sapwood area per individual (cm²/ind).
- `paramsGrowth`: A data frame with growth parameters for each cohort:
 - `WoodC`: Wood carbon content per dry weight (g C /g dry).

- RGRsapwoodmax: Maximum relative growth rate (in basal area or sapwood area) (in $\text{cm}^2 \cdot \text{cm}^{-2}$).
- fHDmin: Minimum value of the height-to-diameter ratio (dimensionless).
- fHDmax: Maximum value of the height-to-diameter ratio (dimensionless).
- paramsAllometry: A data frame with allometric parameters for each cohort:
 - Aash: Regression coefficient relating the square of shrub height with shrub area.
 - Absh, Bbsh: Allometric coefficients relating phytovolume with dry weight of shrub individuals.
 - Acr, B1cr, B2cr, B3cr, C1cr, C2cr: Regression coefficients used to calculate crown ratio of trees.
 - Acw, Bcw: Regression coefficients used to calculate crown width of trees.
- internalAllocation: A data frame with internal allocation variables for each cohort:
 - allocationTarget: Value of the allocation target variable.
 - leafAreaTarget: Target leaf area (m^2) per individual.
 - fineRootBiomassTarget: Target fine root biomass (g dry) per individual (only if transpirationMode = "Sperry").
- internalCarbon and internalRings: data structures to store other internal state variables.

Author(s)

Miquel De Cáceres Ainsa, CTFC

See Also

[resetInputs](#), [spwb](#), [soil](#), [forest](#), [SpParamsMED](#), [defaultSoilParams](#), [plant_ID](#)

Examples

```
#Load example plot plant data
data(exampleforestMED)

#Default species parameterization
data(SpParamsMED)

# Aboveground parameters
above = forest2aboveground(exampleforestMED, SpParamsMED)
above

# Initialize soil with default soil params
examplesoil = soil(defaultSoilParams())

# Rooting depths
Z50 = c(exampleforestMED$treeData$Z50, exampleforestMED$shrubData$Z50)
Z95 = c(exampleforestMED$treeData$Z95, exampleforestMED$shrubData$Z95)

# Initialize control parameters
control = defaultControl()
```

```

# Prepare spwb input
spwbInput(above, Z50, Z95, examplesoil, SpParamsMED, control)

# When starting from an object of class 'forest' the whole process
# can be simplified:
forest2spwbInput(exampleforestMED, examplesoil, SpParamsMED, control)

# Prepare input for complex transpiration mode
control$transpirationMode = "Sperry"
forest2spwbInput(exampleforestMED, examplesoil, SpParamsMED, control)

```

modifySpParams	<i>Modify species/cohort parameters</i>
----------------	---

Description

Routines to modify input species parameters or model input objects

Usage

```

modifySpParams(SpParams, customParams, subsetSpecies = TRUE)
modifyCohortParams(x, customParams)

```

Arguments

x	A model input object of class <code>spwbInput</code> .
SpParams	A species parameter data frame, typically <code>SpParamsMED</code> .
customParams	A data frame with new parameter values (see details).
subsetSpecies	A flag to indicate that the only species mentioned in <code>customParams</code> should be returned.

Details

Data frame `customParams` should have as many rows as species or cohorts and as many columns as parameters to modify. An additional column is needed to identify species/cohorts. In the case of `modifySpParams` the column should be named 'SpIndex' (to match the corresponding column of `SpParamsMED`), while in `modifyCohortParams` the column should be named 'Cohort' and will be matched with the cohort names given by `spwbInput` or `growthInput`.

Value

Function `modifySpParams` returns a modified species parameter data frame. Function `modifyCohortParams` returns a modified `spwbInput` object.

Author(s)

Miquel De Cáceres Ainsa, CTFC

See Also

[spwbInput](#), [SpParamsMED](#)

Examples

```
## TO DO
```

pheno_updateLeaves *Leaf phenology*

Description

Function `pheno_leafDevelopmentStatus` returns the expanded status (0 to 1) of leaves according to the growth degree days required to start bud burst and leaf unfolding, as dictated by a simple ecodormancy (one-phase) model (Chuine et al. 2013). Function `pheno_leafSenescenceStatus` returns the 0/1 senescence status of leaves according to the one-phase senescence model of Delpierre et al. (2009) on the basis of photoperiod and temperature. Function `pheno_updateLeaves` updates the status of expanded leaves and dead leaves of object `x` given the photoperiod, temperature and wind of a given day. It applies the development model for $1 < \text{doy} < 180$ and the senescence model for $181 > \text{doy} > 365$.

Usage

```
pheno_leafDevelopmentStatus(Sgdd, gdd, unfoldingDD = 300)
pheno_leafSenescenceStatus(Ssen, sen)
pheno_updatePhenology(x, doy, photoperiod, tmean)
pheno_updateLeaves(x, wind, fromGrowthModel)
```

Arguments

<code>Sgdd</code>	Degree days required for leaf budburst (in Celsius).
<code>gdd</code>	Cumulative degree days (in Celsius)
<code>unfoldingDD</code>	Degree-days for complete leaf unfolding after budburst has occurred.
<code>Ssen</code>	Threshold to start leaf senescence.
<code>sen</code>	Cumulative senescence variable.
<code>x</code>	An object of class spwbInput .
<code>doy</code>	Day of the year.
<code>photoperiod</code>	Day length (in hours).
<code>tmean</code>	Average day temperature (in Celsius).
<code>wind</code>	Average day wind speed (in m/s).
<code>fromGrowthModel</code>	Boolean flag to indicate that routine is called from growth simulation function.


```

absRadSL, absRadSH, QSL, QSH,
Vmax298SL, Vmax298SH, Jmax298SL, Jmax298SH,
leafWidth = 1.0, verbose = FALSE)
photo_multilayerPhotosynthesisFunction(E, Catm, Patm, Tair, vpa,
SLarea, SHarea, u,
absRadSL, absRadSH, QSL, QSH,
Vmax298, Jmax298, leafWidth = 1.0,
verbose = FALSE)

```

Arguments

Tleaf	Leaf temperature (in °C).
Oi	Oxygen concentration (mmol*mol ⁻¹).
Vmax298, Vmax298SL, Vmax298SH	Maximum Rubisco carboxylation rate per leaf area at 298°K (i.e. 25 °C) (micromol*s ⁻¹ *m ⁻²) (for each canopy layer in the case of photo_multilayerPhotosynthesisFunction). 'SH' stands for shade leaves, whereas 'SL' stands for sunlit leaves.
Jmax298, Jmax298SL, Jmax298SH	Maximum electron transport rate per leaf area at 298°K (i.e. 25 °C) (micromol*s ⁻¹ *m ⁻²) (for each canopy layer in the case of photo_multilayerPhotosynthesisFunction). 'SH' stands for shade leaves, whereas 'SL' stands for sunlit leaves.
Q	Active photon flux density (micromol * s ⁻¹ * m ⁻²).
Ci	CO2 internal concentration (micromol * mol ⁻¹).
GT	CO2 saturation point corrected by temperature (micromol * mol ⁻¹).
Jmax	Maximum electron transport rate per leaf area (micromol*s ⁻¹ *m ⁻²).
Km	$K_m = K_c * (1.0 + (O_i / K_o))$ - Michaelis-Menten term corrected by temperature (in micromol * mol ⁻¹).
Vmax	Maximum Rubisco carboxylation rate per leaf area (micromol*s ⁻¹ *m ⁻²).
Catm	CO2 air concentration (micromol * mol ⁻¹).
Gc	CO2 leaf (stomatal) conductance (mol * s ⁻¹ * m ⁻²).
E	Transpiration flow rate per leaf area (mmol*s ⁻¹ *m ⁻²).
Patm	Atmospheric air pressure (in kPa).
Tair	Air temperature (in °C).
vpa	Vapour pressure deficit (in kPa).
u	Wind speed above the leaf boundary (in m/s) (for each canopy layer in the case of photo_multilayerPhotosynthesisFunction).
absRad	Absorbed long- and short-wave radiation (in W*m ⁻²).
leafWidth	Leaf width (in cm).
refLeafArea	Leaf reference area.
verbose	Boolean flag to indicate console output.
SLarea, SHarea	Leaf area index of sunlit/shade leaves (for each canopy layer in the case of photo_multilayerPhotosynthesisFunction).

absRadSL, absRadSH	Instantaneous absorbed radiation ($W \cdot m^{-2}$) per unit of sunlit/shade leaf area (for each canopy layer in the case of <code>photo_multilayerPhotosynthesisFunction</code>).
QSL, QSH	Active photon flux density ($\mu mol \cdot s^{-1} \cdot m^{-2}$) per unit of sunlit/shade leaf area (for each canopy layer in the case of <code>photo_multilayerPhotosynthesisFunction</code>).

Details

Details of the photosynthesis submodel are given in a vignette.

Value

Values returned for each function are:

- `photo_GammaTemp`: CO₂ compensation concentration ($\mu mol \cdot mol^{-1}$).
- `photo_KmTemp`: Michaelis-Menten coefficients of Rubisco for Carbon ($\mu mol \cdot mol^{-1}$) and Oxygen ($mmol \cdot mol^{-1}$).
- `photo_VmaxTemp`: Temperature correction of V_{max298} .
- `photo_JmaxTemp`: Temperature correction of J_{max298} .
- `photo_electronLimitedPhotosynthesis`: Electron-limited photosynthesis ($\mu mol \cdot s^{-1} \cdot m^{-2}$) following Farquhar et al. (1980).
- `photo_rubiscoLimitedPhotosynthesis`: Rubisco-limited photosynthesis ($\mu mol \cdot s^{-1} \cdot m^{-2}$) following Farquhar et al. (1980).
- `photo_photosynthesis`: Calculates gross photosynthesis ($\mu mol \cdot s^{-1} \cdot m^{-2}$) following (Farquhar et al. (1980) and Collatz et al (1991).
- `photo_leafPhotosynthesisFunction`: Returns a data frame with the following columns:
 - `LeafTemperature`: Leaf temperature ($^{\circ}C$).
 - `LeafVPD`: Leaf vapor pressure deficit (kPa).
 - `LeafCi`: Internal CO₂ concentration ($\mu mol \cdot mol^{-1}$).
 - `WaterVaporConductance`: Leaf vapor conductance ($mol \cdot s^{-1} \cdot m^{-2}$).
 - `GrossPhotosynthesis`: Gross photosynthesis ($\mu mol \cdot s^{-1} \cdot m^{-2}$).
 - `NetPhotosynthesis`: Net photosynthesis, after discounting autotrophic respiration ($\mu mol \cdot s^{-1} \cdot m^{-2}$).
- `photo_sunshadePhotosynthesisFunction`: Returns a data frame with the following columns:
 - `GrossPhotosynthesis`: Gross photosynthesis ($\mu mol \cdot s^{-1} \cdot m^{-2}$).
 - `NetPhotosynthesis`: Net photosynthesis, after discounting autotrophic respiration ($\mu mol \cdot s^{-1} \cdot m^{-2}$).
 - `LeafCiSL`: Sunlit leaf internal CO₂ concentration ($\mu mol \cdot mol^{-1}$).
 - `LeafCiSH`: Shade leaf internal CO₂ concentration ($\mu mol \cdot mol^{-1}$).
 - `LeafTempSL`: Sunlit leaf temperature ($^{\circ}C$).
 - `LeafTempSH`: Shade leaf temperature ($^{\circ}C$).
 - `LeafVPDSL`: Sunlit leaf vapor pressure deficit (kPa).
 - `LeafVPDSH`: Shade leaf vapor pressure deficit (kPa).
- `photo_multilayerPhotosynthesisFunction`: Return a data frame with the following columns:

- GrossPhotosynthesis: Gross photosynthesis ($\text{micromol*s}^{-1}\text{m}^{-2}$).
- NetPhotosynthesis: Net photosynthesis, after discounting autotrophic respiration ($\text{micromol*s}^{-1}\text{m}^{-2}$).

Author(s)

Miquel De Cáceres Ainsa, CTFC

References

- Bernacchi, C. J., E. L. Singsaas, C. Pimentel, A. R. Portis, and S. P. Long. 2001. Improved temperature response functions for models of Rubisco-limited photosynthesis. *Plant, Cell and Environment* 24:253–259.
- Collatz, G. J., J. T. Ball, C. Grivet, and J. A. Berry. 1991. Physiological and environmental regulation of stomatal conductance, photosynthesis and transpiration: a model that includes a laminar boundary layer. *Agricultural and Forest Meteorology* 54:107–136.
- Farquhar, G. D., S. von Caemmerer, and J. A. Berry. 1980. A biochemical model of photosynthetic CO₂ assimilation in leaves of C₃ species. *Planta* 149:78–90.
- Leuning, R. 2002. Temperature dependence of two parameters in a photosynthesis model. *Plant, Cell and Environment* 25:1205–1210.
- Sperry, J. S., M. D. Venturas, W. R. L. Anderegg, M. Mencuccini, D. S. Mackay, Y. Wang, and D. M. Love. 2016. Predicting stomatal responses to the environment from the optimization of photosynthetic gain and hydraulic cost. *Plant Cell and Environment*.

See Also

[hydraulics_supplyFunctionNetwork](#), [biophysics_leafTemperature](#), [spwb](#)

Plant values

Plant description functions

Description

Functions to calculate attributes of plants in a [forest](#) object.

Usage

```

plant_basalArea(x)
plant_largerTreeBasalArea(x)
plant_characterParameter(x, SpParams, parName)
plant_cover(x)
plant_crownBaseHeight(x, SpParams, mode = "MED")
plant_crownLength(x, SpParams, mode = "MED")
plant_crownRatio(x, SpParams, mode = "MED")
plant_density(x, SpParams, mode = "MED")
plant_equilibriumLeafLitter(x, SpParams, AET = 800)

```

```

plant_equilibriumSmallBranchLitter(x, SpParams,
                                   smallBranchDecompositionRate = 0.81)
plant_foliarBiomass(x, SpParams, gdd = NA, mode = "MED")
plant_fuel(x, SpParams, gdd = NA, includeDead = TRUE, mode = "MED")
plant_height(x)
plant_ID(x)
plant_LAI(x, SpParams, gdd = NA, mode = "MED")
plant_parameter(x, SpParams, parName)
plant_phytovolume(x, SpParams)
plant_species(x)
plant_speciesName(x, SpParams)

```

Arguments

x	An object of class <code>forest</code> .
SpParams	A data frame with species parameters (see <code>SpParamsMED</code>).
parName	A string with a parameter name.
mode	Calculation mode, either "MED" or "US".
gdd	Growth degree days (to account for leaf phenology effects).
AET	Actual annual evapotranspiration (in mm).
smallBranchDecompositionRate	Decomposition rate of small branches.
includeDead	A flag to indicate that standing dead fuels (dead branches) are included.

Value

A vector with values for each plant of the input `forest` object:

- `plant_basalArea`: Tree basal area (m²/ha).
- `plant_largerTreeBasalArea`: Basal area (m²/ha) of trees larger (in diameter) than the tree. Half of the trees of the same record are included.
- `plant_characterParameter`: The parameter values of each plant, as strings.
- `plant_cover`: Shrub cover (in percent).
- `plant_crownBaseHeight`: The height corresponding to the start of the crown (in cm).
- `plant_crownLength`: The difference between crown base height and total height (in cm).
- `plant_crownRatio`: The ratio between crown length and total height (between 0 and 1).
- `plant_density`: Plant density (ind/ha). Tree density is directly taken from the forest object, while the shrub density is estimated from cover and height by calculating the area of a single individual.
- `plant_equilibriumLeafLitter`: Litter biomass of leaves at equilibrium (in kg/m²).
- `plant_equilibriumSmallBranchLitter`: Litter biomass of small branches (< 6.35 mm diameter) at equilibrium (in kg/m²).
- `plant_foliarBiomass`: Standing biomass of leaves (in kg/m²).
- `plant_fuel`: Fine fuel load (in kg/m²).

- `plant_height`: Total height (in cm).
- `plant_ID`: Cohort coding for simulation functions (concatenation of 'T' (Trees) or 'S' (Shrub), cohort index and species index).
- `plant_LAI`: Leaf area index (m²/m²).
- `plant_parameter`: The parameter values of each plant, as numeric.
- `plant_phytovolume`: Shrub phytovolume (m³/m²).
- `plant_species`: Species identity integer (indices start with 0).
- `plant_speciesName`: String with species taxonomic name (or a functional group).

Author(s)

Miquel De Cáceres Ainsa, CTFC

See Also

[spwb](#), [forest](#), [summary.forest](#)

Examples

```
#Default species parameterization
data(SpParamsMED)

#Load example plot
data(exampleforestMED)

#A short way to obtain total basal area
sum(plant_basalArea(exampleforestMED), na.rm=TRUE)

#The same forest level function for LAI
sum(plant_LAI(exampleforestMED, SpParamsMED))

#The same forest level function for fuel loading
sum(plant_fuel(exampleforestMED, SpParamsMED))

#Summary function for 'forest' objects can be also used
summary(exampleforestMED, SpParamsMED)

plant_speciesName(exampleforestMED, SpParamsMED)

plant_ID(exampleforestMED)
```

Description

Function plot produces temporal graphs of the results of the soil plant water balance model (see [spwb](#)), plant water balance model (see [pwb](#)) or the forest growth model (see [growth](#)), whereas function summary summarizes the model's output in different temporal steps (i.e. weekly, annual, ...).

Usage

```
## S3 method for class 'spwb'
plot(x, type="PET_Precipitation",
      cohorts = NULL, bySpecies = FALSE,
      dates = NULL, subdaily = FALSE,
      xlim = NULL, ylim=NULL, xlab=NULL, ylab=NULL, ...)
## S3 method for class 'pwb'
plot(x, type="PlantTranspiration",
      cohorts = NULL, bySpecies = FALSE,
      dates = NULL, subdaily = FALSE,
      xlim = NULL, ylim=NULL, xlab=NULL, ylab=NULL, ...)
## S3 method for class 'growth'
plot(x, type="PET_Precipitation",
      cohorts = NULL, bySpecies = FALSE,
      dates = NULL, subdaily = FALSE,
      xlim = NULL, ylim=NULL, xlab=NULL, ylab=NULL, ...)
## S3 method for class 'pwb'
summary(object, freq="years", output="WaterBalance", FUN=sum, bySpecies = FALSE, ...)
## S3 method for class 'spwb'
summary(object, freq="years", output="WaterBalance", FUN=sum, bySpecies = FALSE, ...)
## S3 method for class 'growth'
summary(object, freq="years", output="WaterBalance", FUN=sum, bySpecies = FALSE, ...)
```

Arguments

x, object	An object of class spwb.
type	The information to be plotted (see details)
cohorts	An integer, boolean or character vector to select the plant cohorts to be plotted.
bySpecies	Allows aggregating output by species, before calculating summaries or drawing plots (only has an effect with some values of type). Aggregation can involve a sum (as for plant lai or transpiration) or a LAI-weighted mean (as for plant stress or plant water potential).
dates	A Date vector with a subset of dates to be plotted.
subdaily	Whether subdaily results should be shown, only for simulations using transpirationMode = "Sperry" and having set subdailyResults = TRUE in the simulation control object. If subdaily = TRUE, then the valid strings for type are listed in plot.spwb_day .
xlim	Range of values for x.
ylim	Range of values for y.

xlab	x-axis label.
ylab	y-axis label.
freq	Frequency of summary statistics (see cut.Date).
output	The data table to be summarized. Accepted values are "DailyBalance", "PlantStress", "PlantPsi", "PlantTranspiration", "PlantPhotosynthesis" and "SoilWaterBalance", "Temperature" and "EnergyBalance".
FUN	The function to summarize results (e.g., sum, mean, ...)
...	Additional parameters for function plot or summary.

Details

The following plots are currently available for [spwb](#) (most of them also for [pwb](#)):

- "PET_Precipitation": Potential evapotranspiration and Precipitation.
- "PET_NetRain": Potential evapotranspiration and Net rainfall.
- "Snow": Snow precipitation and snowpack dynamics.
- "Export": Water exported through deep drainage and surface runoff.
- "Evapotranspiration": Plant transpiration and soil evaporation.
- "SoilPsi": Soil water potential.
- "SoilRWC": Soil relative water content (in percent of field capacity).
- "SoilTheta": Soil moisture water content (in percent volume).
- "SoilVol": Soil water volumetric content (in mm).
- "PlantExtraction": Water extracted by plants from each soil layer.
- "HydraulicRedistribution": Water added to each soil layer coming from other soil layers, transported through the plant hydraulic network (only for transpirationMode = "Sperry").
- "WTD": Water table depth.
- "LAI": Expanded and dead leaf area index of the whole stand.
- "PlantLAI": Plant cohort leaf area index (expanded leaves).
- "SoilPlantConductance": Average instantaneous overall soil plant conductance (calculated as the derivative of the supply function).
- "PlantStress": Plant cohort average daily drought stress.
- "PlantPsi": Plant cohort water potential (only for transpirationMode = "Granier").
- "LeafPsi": Midday leaf water potential (only for transpirationMode = "Sperry").
- "StemPsi": Midday (upper) stem water potential (only for transpirationMode = "Sperry").
- "RootPsi": Midday root crown water potential (only for transpirationMode = "Sperry").
- "PlantTranspiration": Plant cohort transpiration.
- "PlantTranspirationPerLeaf": Plant cohort transpiration per leaf area.
- "PlantPhotosynthesis": Plant cohort photosynthesis.
- "PlantPhotosynthesisPerLeaf": Plant cohort photosynthesis per leaf area.
- "PlantWUE": Plant cohort daily water use efficiency (photosynthesis over transpiration).

- "PlantAbsorbedSWR": Plant cohort absorbed short wave radiation (only for transpirationMode = "Sperry").
- "PlantAbsorbedSWRPerLeaf": Plant cohort absorbed short wave radiation per leaf area (only for transpirationMode = "Sperry").
- "PlantAbsorbedLWR": Plant cohort absorbed long wave radiation (only for transpirationMode = "Sperry").
- "PlantAbsorbedLWRPerLeaf": Plant cohort absorbed long wave radiation per leaf area (only for transpirationMode = "Sperry").
- "AirTemperature": Minimum/maximum/mean daily temperatures above canopy (only for transpirationMode = "Sperry").
- "CanopyTemperature": Minimum/maximum/mean daily temperatures inside canopy (only for transpirationMode = "Sperry").
- "SoilTemperature": Minimum/maximum/mean daily temperatures inside the first soil layer (only for transpirationMode = "Sperry").
- "CanopyEnergyBalance": Canopy energy balance components (only for transpirationMode = "Sperry").
- "SoilEnergyBalance": Soil energy balance components (only for transpirationMode = "Sperry").

The following are only available for [growth](#):

- "PlantRespiration": Plant cohort respiration.
- "PlantRespirationPerLeaf": Plant cohort respiration per leaf area.
- "PlantRespirationPerIndividual": Respiration per individual.
- "PlantCBalance": Plant cohort carbon balance.
- "PlantCBalancePerLeaf": Plant cohort carbon balance per leaf area.
- "PlantCBalancePerIndividual": Carbon balance per individual.
- "PlantCstorageFast": Amount of fast-dynamics carbon reserves.
- "PlantCstorageSlow": Amount of slow-dynamics carbon reserves.
- "PlantSA": Amount of sapwood area in an individual.
- "PlantSAGrowth": Amount of newly-created sapwood area.
- "PlantRelativeSAGrowth": Amount of newly-created sapwood area per sapwood area.
- "PlantLAIlive": Plant cohort leaf area index of live leaves.
- "PlantLAIdead": Plant cohort leaf area index of dead leaves.

Author(s)

Miquel De Cáceres Ainsa, CTFC

References

De Cáceres M, Martínez-Vilalta J, Coll L, Llorens P, Casals P, Poyatos R, Pausas JG, Brotons L. (2015) Coupling a water balance model with forest inventory data to predict drought stress: the role of forest structural changes vs. climate changes. *Agricultural and Forest Meteorology* 213: 77-90 (doi:10.1016/j.agrformet.2015.06.012).

See Also[spwb](#)**Examples**

```

#Load example daily meteorological data
data(examplemeteo)

#Load example plot plant data
data(exampleforestMED)

#Default species parameterization
data(SpParamsMED)

#Initialize soil with default soil params (2 layers)
examplesoil = soil(defaultSoilParams(2))

#Initialize control parameters
control = defaultControl()

#Initialize input
x = forest2spwbInput(exampleforestMED,examplesoil, SpParamsMED, control)

#Call simulation function
S1<-spwb(x, examplesoil, examplemeteo, latitude = 41.82592, elevation = 100)

#Plot results
plot(S1)

#Monthly summary (averages) of soil status
summary(S1, freq="months",FUN=mean, output="Soil")

```

plot.spwb_day

Plots simulation results for one day

Description

Functions to plot the subdaily simulation results of [spwb_day](#), [growth_day](#) or the transpiration calculations of [transp_transpirationSperry](#).

Usage

```

## S3 method for class 'growth_day'
plot(x, type="PlantTranspiration", bySpecies = FALSE,
      xlim = NULL, ylim=NULL, xlab = NULL, ylab = NULL, ...)
## S3 method for class 'spwb_day'
plot(x, type="PlantTranspiration", bySpecies = FALSE,
      xlim = NULL, ylim=NULL, xlab = NULL, ylab = NULL, ...)

```

```
## S3 method for class 'pwb_day'
plot(x, type="PlantTranspiration", bySpecies = FALSE,
      xlim = NULL, ylim=NULL, xlab = NULL, ylab = NULL, ...)
```

Arguments

x	An object of class spwb_day, growth_day or pwb_day.
type	The information to be plotted (see details).
bySpecies	Allows aggregating output by species, before drawing plots. Aggregation can involve a sum (as for plant lai or transpiration) or a LAI-weighted mean (as for plant stress or plant water potential).
xlim	Range of values for x.
ylim	Range of values for y.
xlab	x-axis label.
ylab	y-axis label.
...	Additional parameters for function plot.

Details

The following plots are currently available for spwb_day and pwb_day:

- "LeafPsi": Leaf water potential (for shade and sunlit leaves).
- "LeafPsiAverage": Average leaf water potential.
- "RootPsi": Root crown water potential.
- "StemPsi": (Upper) stem water potential.
- "StemPLC": (Average) percentage of loss conductance in the stem conduits.
- "StemRWC": (Average) relative water content in the stem.
- "LeafRWC": Relative water content in the leaf.
- "StemSympRWC": (Average) relative water content in the stem symplasm.
- "LeafSympRWC": Relative water content in the leaf symplasm.
- "SoilPlantConductance": Overall soil plant conductance (calculated as the derivative of the supply function).
- "PlantExtraction": Water extracted from each soil layer.
- "PlantTranspiration": Plant cohort transpiration per ground area.
- "TranspirationPerLeaf": Plant cohort transpiration per leaf area.
- "PlantGrossPhotosynthesis": Plant cohort gross photosynthesis per ground area.
- "GrossPhotosynthesisPerLeaf": Plant cohort gross photosynthesis per leaf area.
- "PlantNetPhotosynthesis": Plant cohort net photosynthesis per ground area.
- "NetPhotosynthesisPerLeaf": Plant cohort net photosynthesis per leaf area.
- "PlantAbsorbedSWR": Absorbed short wave radiation per ground area (differentiates sunlit and shade leaves).

- "LeafTranspiration": Instantaneous transpiration per leaf area (differentiates sunlit and shade leaves).
- "LeafGrossPhotosynthesis": Instantaneous gross photosynthesis per leaf area (differentiates sunlit and shade leaves).
- "LeafNetPhotosynthesis": Instantaneous net photosynthesis per leaf area (differentiates sunlit and shade leaves).
- "LeafAbsorbedSWR": Absorbed short wave radiation per leaf area (differentiates sunlit and shade leaves).
- "LeafAbsorbedLWR": Absorbed long wave radiation per leaf area (differentiates sunlit and shade leaves).
- "LeafCi": Leaf intercellular CO₂ concentration (differentiates sunlit and shade leaves).
- "LeafIntrinsicWUE": Leaf intrinsic water use efficiency, i.e. the ratio between instantaneous photosynthesis and stomatal conductance (differentiates sunlit and shade leaves).
- "LeafVPD": Leaf vapour pressure deficit (differentiates sunlit and shade leaves).
- "LeafStomatalConductance": Leaf stomatal conductance to water vapour (differentiates sunlit and shade leaves).
- "LeafTemperature": Leaf temperature (differentiates sunlit and shade leaves).
- "Temperature": Above-canopy, inside-canopy and soil temperature.
- "CanopyEnergyBalance": Canopy energy balance components.
- "SoilEnergyBalance": Soil energy balance components.
- "PlantWaterBalance": Difference between water extraction from the soil and transpired water per ground area.
- "WaterBalancePerLeaf": Difference between water extraction from the soil and transpired water per leaf area.

Note

Only for soil plant water balance simulations using `transpirationMode = "Sperry"`. This function can be used to display subdaily dynamics of corresponding to single days on `spwb` runs, if control option `subdailyResults` is set to `TRUE`. See also option `subdaily` in `plot.spwb`.

Author(s)

Miquel De Cáceres Ainsa, CTFC

See Also

`spwb_day`, `plot.spwb`

Examples

```
#Load example daily meteorological data
data(examplemeteo)

#Load example plot plant data
```

```

data(exampleforestMED)

#Default species parameterization
data(SpParamsMED)

#Initialize control parameters
control = defaultControl()
control$ndailysteps = 24

#Initialize soil with default soil params (2 layers)
examplesoil = soil(defaultSoilParams(2), W=c(0.5,0.5))

#Switch to 'Sperry' transpiration mode
control$transpirationMode="Sperry"

#Simulate one day only
x2 = forest2spwbInput(exampleforestMED,examplesoil, SpParamsMED, control)
d = 100
sd2<-spwb_day(x2, examplesoil, rownames(examplemeteo)[d],
              examplemeteo$MinTemperature[d], examplemeteo$MaxTemperature[d],
              examplemeteo$MinRelativeHumidity[d], examplemeteo$MaxRelativeHumidity[d],
              examplemeteo$Radiation[d], examplemeteo$WindSpeed[d],
              latitude = 41.82592, elevation = 100,
              slope= 0, aspect = 0, prec = examplemeteo$Precipitation[d])

#Display transpiration for subdaily steps
plot(sd2, "PlantTranspiration")

```

resetInputs

Reset simulation inputs

Description

Function `resetInputs()` allows resetting state variables in `x` and `soil` to their defaults.

Usage

```
resetInputs(x, soil)
```

Arguments

`x` An object of class `spwbInput` or `growthInput`.

`soil` A list containing the description of the soil (see `soil`).

Value

Does not return any value. Instead, it modifies input objects `x` and `soil`.

Author(s)

Miquel De Cáceres Ainsa, CTFC

See Also

[spwbInput](#), [growthInput](#), [spwb](#)

root

Root functions

Description

Functions to calculate properties of fine/coarse roots within the soil, given root system parameters and soil layer definition.

Usage

```

root_conicDistribution(Zcone, d)
root_ldrDistribution(Z50, Z95, d)
root_coarseRootLengths(v, d, depthWidthRatio = 1.0)
root_coarseRootLengthsFromVolume(VolInd, v, d, rfc)
root_coarseRootSoilVolume(v, d, depthWidthRatio = 1.0)
root_coarseRootSoilVolumeFromConductance(Kmax_rootxylem, VCrout_kmax, A12As,
                                          v, d, rfc)
root_fineRootRadius(specificRootLength, rootTissueDensity)
root_fineRootHalfDistance(rootLengthDensity)
root_fineRootAreaIndex(Ksoil, krhizo, lai,
                      specificRootLength, rootTissueDensity,
                      rootLengthDensity)
root_fineRootBiomass(Ksoil, krhizo, lai, N,
                   specificRootLength, rootTissueDensity, rootLengthDensity)
root_rhizosphereMaximumConductance(Ksoil, fineRootBiomass, lai, N,
                                   specificRootLength, rootTissueDensity,
                                   rootLengthDensity)
root_fineRootSoilVolume(fineRootBiomass, specificRootLength, rootLengthDensity)
root_specificRootSurfaceArea(specificRootLength, rootTissueDensity)
root_individualRootedGroundArea(VolInd, V, d, rfc)
root_horizontalProportions(poolProportions, VolInd, N, V, d, rfc)

```

Arguments

Z50	A vector of depths (in mm) corresponding to 50% of roots.
Z95	A vector of depths (in mm) corresponding to 95% of roots.
Zcone	A vector of depths (in mm) corresponding to the root cone tip.
d	The width (in mm) corresponding to each soil layer.
v	Vector of proportions of fine roots in each soil layer.

depthWidthRatio	Ratio between radius of the soil layer with the largest radius and maximum rooting depth.
rfc	Percentage of rock fragment content (volume basis) for each layer.
Kmax_rootxylem	Sapwood-specific hydraulic conductivity of root xylem (in $\text{kg H}_2\text{O}\cdot\text{s}^{-1}\cdot\text{m}^{-1}\cdot\text{MPa}^{-1}$).
VCroot_kmax	Root xylem maximum conductance per leaf area ($\text{mmol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}\cdot\text{MPa}^{-1}$).
Al2As	Leaf area to sapwood area ratio (in $\text{m}^2\cdot\text{m}^{-2}$).
specificRootLength	Specific fine root length (length of fine roots over weight).
rootTissueDensity	Fine root tissue density (weight over volume at turgidity).
Ksoil	Soil saturated conductivity ($\text{mmol}\cdot\text{m}^{-1}\cdot\text{s}^{-1}\cdot\text{MPa}^{-1}$).
krhizo	Rhizosphere maximum conductance per leaf area ($\text{mmol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}\cdot\text{MPa}^{-1}$).
lai	Leaf area index.
rootLengthDensity	Fine root length density (length of fine roots over soil volume; cm/cm^3)
fineRootBiomass	Biomass of fine roots (g).
V	Matrix of proportions of fine roots (cohorts x soil layers).
VolInd	Volume of soil (in m^3) occupied by coarse roots per individual.
N	Density of individuals per hectare.
poolProportions	Division of the stand area among plant cohorts (proportions).

Details

- `root_conicDistribution()` assumes a (vertical) conic distribution of fine roots, whereas `root_ldrDistribution()` distributes fine roots according to the linear dose response model of Schenck & Jackson (2002). Return a matrix of fine root proportions in each layer with as many rows as elements in Z (or Z50) and as many columns as soil layers.
- `root_coarseRootLengths()` and `root_coarseRootLengthsFromVolume()` estimate the length of coarse roots (mm) for each soil layer, including axial and radial lengths.
- `root_coarseRootSoilVolume` estimates the soil volume (m^3) occupied by coarse roots of an individual.
- `root_coarseRootSoilVolumeFromConductance` estimates the soil volume (m^3) occupied by coarse roots of an individual from root xylem conductance.
- `root_fineRootHalfDistance()` calculates the half distance (cm) between neighbouring fine roots.
- `root_fineRootRadius()` calculates the radius of fine roots (cm).
- `root_fineRootAreaIndex()` estimates the fine root area index for a given soil conductivity and maximum rhizosphere conductance.

- `root_fineRootBiomass()` estimates the biomass of fine roots (g dry/individual) for a given soil conductivity and maximum rhizosphere conductance.
- `root_rhizosphereMaximumConductance()` is the inverse of the preceding function, i.e. it estimates rhizosphere conductance from soil conductivity and fine root biomass.
- `root_fineRootSoilVolume()` calculates the soil volume (m³) occupied with fine roots.
- `root_specificRootSurfaceArea()` returns the specific fine root area (cm²/g).
- `root_individualRootedGroundArea()` calculates the area (m²) covered by roots of an individual, for each soil layer.
- `root_horizontalProportions()` calculates the (horizontal) proportion of roots of each cohort in the water pool corresponding to itself and that of other cohorts, for each soil layer. Returns a list (with as many elements as cohorts) with each element being a matrix.

Author(s)

Miquel De Cáceres Ainsa, CTFC

References

- Schenk, H., Jackson, R., 2002. The global biogeography of roots. *Ecol. Monogr.* 72, 311–328.
- Sperry, J. S., Y. Wang, B. T. Wolfe, D. S. Mackay, W. R. L. Anderegg, N. G. McDowell, and W. T. Pockman. 2016. Pragmatic hydraulic theory predicts stomatal responses to climatic water deficits. *New Phytologist* 212, 577–589.

See Also

[spwb](#), [spwb_ldrOptimization](#), [forest2spwbInput](#), [soil](#)

Examples

```
#Load example plot plant data
data(exampleforestMED)

#Default species parameterization
data(SpParamsMED)

ntree = nrow(exampleforestMED$treeData)

#Initialize soil with default soil params
S = soil(defaultSoilParams())

#Calculate conic root system for trees
V1 = root_conicDistribution(Z=rep(2000,ntree), S$dVec)
print(V1)

#Calculate LDR root system for trees (Schenck & Jackson 2002)
V2 = root_ldrDistribution(Z50 = rep(200,ntree),
                        Z95 = rep(1000,ntree), S$dVec)
print(V2)
```

scalingconductance *Scaling from conductivity to conductance*

Description

Functions used to scale from tissue conductivity to conductance of different elements of the continuum.

Usage

```

hydraulics_maximumSoilPlantConductance(krhizomax, krootmax,
                                         kstemmax, kleafmax)
hydraulics_soilPlantResistances(psiSoil, psiRhizo,
                                psiStem, PLCstem, psiLeaf,
                                krhizomax, n, alpha,
                                krootmax, rootc, rootd,
                                kstemmax, stemc, stemd,
                                kleafmax, leafc, leafd)
hydraulics_averageRhizosphereResistancePercent(krhizomax, n, alpha,
                                                krootmax, rootc, rootd,
                                                kstemmax, stemc, stemd,
                                                kleafmax, leafc, leafd, psiStep = -0.01)
hydraulics_findRhizosphereMaximumConductance(averageResistancePercent, n, alpha,
                                                krootmax, rootc, rootd,
                                                kstemmax, stemc, stemd,
                                                kleafmax, leafc, leafd, initialValue = 0)
hydraulics_maximumStemHydraulicConductance(xylemConductivity, refheight, Al2As, height,
                                             taper = FALSE)
hydraulics_rootxylemConductanceProportions(L, V)
hydraulics_referenceConductivityHeightFactor(refheight, height)
hydraulics_terminalConduitRadius(height)
hydraulics_taperFactorSavage(height)
hydraulics_stemWaterCapacity(Al2As, height, wd)
hydraulics_leafWaterCapacity(SLA, ld)

```

Arguments

psiSoil	Soil water potential (in MPa). A scalar or a vector depending on the function.
psiRhizo	Water potential (in MPa) in the rhizosphere (root surface).
psiStem	Water potential (in MPa) in the stem.
psiLeaf	Water potential (in MPa) in the leaf.
PLCstem	Percent loss of conductance (in %) in the stem.
L	Vector with the length of coarse roots (mm) for each soil layer.
V	Vector with the proportion [0-1] of fine roots within each soil layer.

krhizomax	Maximum rhizosphere hydraulic conductance (defined as flow per leaf surface unit and per pressure drop).
kleafmax	Maximum leaf hydraulic conductance (defined as flow per leaf surface unit and per pressure drop).
kstemmax	Maximum stem xylem hydraulic conductance (defined as flow per leaf surface unit and per pressure drop).
krootmax	Maximum root xylem hydraulic conductance (defined as flow per leaf surface unit and per pressure drop).
psiStep	Water potential precision (in MPa).
rootc, rootd	Parameters of the Weibull function for roots (root xylem vulnerability curve).
stemc, stemd	Parameters of the Weibull function for stems (stem xylem vulnerability curve).
leafc, leafd	Parameters of the Weibull function for leaves (leaf vulnerability curve).
n, alpha	Parameters of the Van Genuchten function (rhizosphere vulnerability curve).
averageResistancePercent	Average (across water potential values) resistance percent of the rhizosphere, with respect to total resistance (rhizosphere + root xylem + stem xylem).
initialValue	Initial value of rhizosphere conductance.
xylemConductivity	Xylem conductivity as flow per length of conduit and pressure drop (in $\text{kg}\cdot\text{m}^{-1}\cdot\text{s}^{-1}\cdot\text{MPa}^{-1}$).
A12As	Leaf area to sapwood area (in $\text{m}^2\cdot\text{m}^{-2}$).
height	Plant height (in cm).
refheight	Reference plant height of measurement of xylem conductivity (in cm).
taper	A boolean flag to indicate correction by taper of xylem conduits (Christoffersen et al. 2017).
SLA	Specific leaf area ($\text{mm}^2\cdot\text{mg}^{-1}$).
wd	Wood density ($\text{g}\cdot\text{cm}^{-3}$).
ld	Leaf tissue density ($\text{g}\cdot\text{cm}^{-3}$).

Details

Details of the hydraulic model are given in a vignette.

Value

Values returned for each function are:

- `hydraulics_maximumSoilPlantConductance`: The maximum soil-plant conductance, in the same units as the input segment conductances.
- `hydraulics_averageRhizosphereResistancePercent`: The average percentage of resistance due to the rhizosphere, calculated across water potential values.
- `hydraulics_findRhizosphereMaximumConductance`: The maximum rhizosphere conductance value given an average rhizosphere resistance and the vulnerability curves of rhizosphere, root and stem elements.
- `hydraulics_taperFactorSavage`: Taper factor according to Savage et al. (2010).

Author(s)

Miquel De Cáceres Ainsa, CTFC

References

Christoffersen, B. O., M. Gloor, S. Fauset, N. M. Fyllas, D. R. Galbraith, T. R. Baker, L. Rowland, R. A. Fisher, O. J. Binks, S. A. Sevanto, C. Xu, S. Jansen, B. Choat, M. Mencuccini, N. G. McDowell, and P. Meir. 2016. Linking hydraulic traits to tropical forest function in a size-structured and trait-driven model (TFS v.1-Hydro). *Geoscientific Model Development Discussions* 9: 4227–4255.

Savage, V. M., L. P. Bentley, B. J. Enquist, J. S. Sperry, D. D. Smith, P. B. Reich, and E. I. von Allmen. 2010. Hydraulic trade-offs and space filling enable better predictions of vascular structure and function in plants. *Proceedings of the National Academy of Sciences of the United States of America* 107:22722–7.

Olson, M.E., Anfodillo, T., Rosell, J.A., Petit, G., Crivellaro, A., Isnard, S., León-Gómez, C., Alvarado-Cárdenas, L.O., & Castorena, M. 2014. Universal hydraulics of the flowering plants: Vessel diameter scales with stem length across angiosperm lineages, habits and climates. *Ecology Letters* 17: 988–997.

See Also

[hydraulics_psi2K](#), [hydraulics_supplyFunctionPlot](#), [spwb](#), [soil](#)

Examples

```
kstemmax = 4 # in mmol·m-2·s-1·MPa-1
stemc = 3
stemd = -4 # in MPa
```

SFM_metric

Standard fuel models (Albini 1976, Scott & Burgan 2005)

Description

Standard fuel models converted to metric system. Copied from package 'Rothermel' (Giorgio Vacchiano, Davide Ascoli).

Usage

```
data("SFM_metric")
```

Format

A data frame including standard fuel models as in Albini (1976) and Scott and Burgan (2005), to be used as input of [fire_Rothermel](#) function. All values converted to metric format.

Fuel_Model_Type A factor with levels D (for dynamic) or S (for static).

Load_1h Loading of 1h fuel class [t/ha].

Load_10h Loading of 10h fuel class [t/ha].

Load_100h Loading of 100h fuel class [t/ha]

Load_Live_Herb Loading of herbaceous fuels [t/ha]

Load_Live_Woody Loading of woody fuels [t/ha]

‘SA/V_1h’ Surface area to volume ratio of 1h fuel class [m²/m³]

‘SA/V_10h’ Surface area to volume ratio of 10h fuel class [m²/m³]

‘SA/V_100h’ Surface area to volume ratio of 100h fuel class [m²/m³]

‘SA/V_Live_Herb’ Surface area to volume ratio of herbaceous fuels [m²/m³]

‘SA/V_Live_Woody’ Surface area to volume ratio of woody fuels [m²/m³]

Fuel_Bed_Depth Fuel bed depth [cm]

Mx_dead Dead fuel moisture of extinction [percent]

Heat_1h Heat content of 1h fuel class [kJ/kg]

Heat_10h Heat content of 10h fuel class [kJ/kg]

Heat_100h Heat content of 100h fuel class [kJ/kg]

Heat_Live_Herb Heat content of herbaceous fuels [kJ/kg]

Heat_Live_Woody Heat content of woody fuels [kJ/kg]

Source

Albini, F. A. (1976). Computer-based models of wildland fire behavior: A users’ manual. Ogden, UT: US Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station.

Scott, J., & Burgan, R. E. (2005). A new set of standard fire behavior fuel models for use with Rothermel’s surface fire spread model. Gen. Tech. Rep. RMRS-GTR-153. Fort Collins, CO: US Department of Agriculture, Forest Service, Rocky Mountain Research Station.

See Also

[fire_Rothermel](#)

Examples

```
data(SFM_metric)
```

soil	<i>Soil initialization</i>
------	----------------------------

Description

Initializes soil parameters and state variables for its use in simulations.

Usage

```
soil(SoilParams, VG_PTF = "Toth", W = as.numeric(c(1)), SWE = 0)
## S3 method for class 'soil'
print(x, model="SX", ...)
redefineSoilLayers(SoilParams, widths = c(300, 700, 1000, 2000))
```

Arguments

SoilParams	A data frame of soil parameters (see an example in defaultSoilParams).
VG_PTF	Pedotransfer functions to obtain parameters for the van Genuchten-Mualem equations. Either "Carsel" (Carsel & Parrish 1988) or "Toth" (Toth et al. 2015).
W	A numerical vector with the initial relative water content of each soil layer.
SWE	Initial snow water equivalent of the snow pack on the soil surface (mm).
x	An object of class soil.
model	Either 'SX' or 'VG' for Saxton or Van Genuchten pedotransfer models.
widths	A numeric vector indicating the desired layer widths, in <i>mm</i> .
...	Additional parameters to print.

Details

Function `print` prompts a description of soil characteristics and state variables (water content and temperature) according to a water retention curve (either Saxton's or Van Genuchten's). Volume at field capacity is calculated assuming a soil water potential equal to -0.033 MPa. Parameter `Temp` is initialized as missing for all soil layers. Function `redefineSoilLayers` allows redefining soil layer widths of an input data frame of soil parameters.

Value

Function `soil` returns a list of class `soil` with the following elements:

- `SoilDepth`: Soil depth (in mm).
- `W`: State variable with relative water content of each layer (in as proportion relative to FC).
- `Temp`: State variable with temperature (in °C) of each layer.
- `Ksoil`: Kappa parameter for infiltration.
- `Gsoil`: Gamma parameter for infiltration.
- `dVec`: Width of soil layers (in mm).

- sand: Sand percentage for each layer (in percent volume).
- clay: Clay percentage for each layer (in percent volume).
- om: Organic matter percentage for each layer (in percent volume).
- VG_alpha, VG_n, VG_theta_res, VG_theta_sat: Parameters for van Genuchten's pedotransfer functions, for each layer, corresponding to the USDA texture type.
- Ksat: Saturated soil conductivity for each layer (estimated using function [soil_saturatedConductivitySX](#)).
- macro: Macroporosity for each layer (estimated using Stolf et al. 2011).
- rfc: Percentage of rock fragment content for each layer.
- Kdrain: Saturated vertical hydraulic conductivity (mm/day) (i.e. how easy is deep drainage towards groundwater). Function `soil` estimates it as a function of soil saturated hydraulic conductivity, but should be parametrized as a function of bedrock material.

Author(s)

Miquel De Cáceres Ainsa, CTFC

References

- Carsel, R.F., & Parrish, R.S. 1988. Developing joint probability distributions of soil water retention characteristics. *Water Resources Research* 24: 755–769.
- Tóth, B., Weynants, M., Nemes, A., Makó, A., Bilas, G., & Tóth, G. 2015. New generation of hydraulic pedotransfer functions for Europe. *European Journal of Soil Science* 66: 226–238.
- Stolf, R., Thurler, A., Oliveira, O., Bacchi, S., Reichardt, K., 2011. Method to estimate soil macroporosity and microporosity based on sand content and bulk density. *Rev. Bras. Ciências do Solo* 35, 447–459.

See Also

[soil_psi2thetaSX](#), [soil_psi2thetaVG](#), [spwb](#), [defaultSoilParams](#)

Examples

```
# Initializes soil
s = soil(defaultSoilParams())

# Prints soil characteristics according to Saxton's water retention curve
print(s, model="SX")

# Prints soil characteristics according to Van Genuchten's water retention curve
print(s, model="VG")
```

Description

Function `hydrology_infiltrationAmount` calculates the amount of water that infiltrates into the topsoil, according to the USDA SCS curve number method (Boughton 1989). The remaining is assumed to be lost as surface runoff. Function `hydrology_soilEvaporationAmount` calculates the amount of evaporation from bare soil, following Ritchie (1972). Function `hydrology_snowMelt` calculates the maximum amount of snowmelt according to Kergoat (1998).

Usage

```
hydrology_infiltrationAmount(input, Ssoil)
hydrology_infiltrationRepartition(I, dVec, macro, a = -0.005, b = 3)
hydrology_soilEvaporationAmount(DEF, PETs, Gsoil)
hydrology_soilEvaporation(soil, soilFunctions, pet, LgroundSWR, modifySoil = TRUE)
hydrology_snowMelt(tday, rad, LgroundSWR, elevation)
```

Arguments

<code>input</code>	A numeric vector of (daily) water input (in mm of water).
<code>Ssoil</code>	Soil water storage capacity (can be referred to topsoil) (in mm of water).
<code>DEF</code>	Water deficit in the (topsoil) layer.
<code>PETs</code>	Potential evapotranspiration at the soil surface.
<code>Gsoil</code>	Gamma parameter (maximum daily evaporation).
<code>I</code>	Soil infiltration (in mm of water).
<code>dVec</code>	Width of soil layers (in mm).
<code>macro</code>	Macroporosity of soil layers (in %).
<code>soil</code>	An object of class <code>soil</code> .
<code>soilFunctions</code>	Soil water retention curve and conductivity functions, either 'SX' (for Saxton) or 'VG' (for Van Genuchten).
<code>pet</code>	Potential evapotranspiration for a given day (mm)
<code>LgroundSWR</code>	Proportion of short-wave radiation (SWR) reaching the ground.
<code>modifySoil</code>	Boolean flag to indicate that the input soil object should be modified during the simulation.
<code>tday</code>	Average day temperature (°C).
<code>rad</code>	Solar radiation (in MJ/m ² /day).
<code>elevation</code>	Altitude above sea level (m).
<code>a,b</code>	Parameters of the extinction function used for water infiltration.

Details

See description of infiltration and soil evaporation processes in De Cáceres et al. (2015).

Value

Function `hydrology_infiltrationAmount` a vector of the same length as input containing the daily amount of water that infiltrates into the soil (in mm of water). Function `hydrology_infiltrationRepartition` estimates the amount of infiltrated water that reaches each soil layer. Function `hydrology_soilEvaporationAmount` returns the amount of water evaporated from the soil. Function `hydrology_soilEvaporation` returns a vector of water evaporated from each soil layer.

Author(s)

Miquel De Cáceres Ainsa, CTFC

References

Boughton (1989). A review of the USDA SCS curve number method. - Australian Journal of Soil Research 27: 511-523.

De Cáceres M, Martínez-Vilalta J, Coll L, Llorens P, Casals P, Poyatos R, Pausas JG, Brotons L. (2015) Coupling a water balance model with forest inventory data to evaluate plant drought stress at the regional level. Agricultural and Forest Meteorology.

Kergoat L. (1998). A model for hydrological equilibrium of leaf area index on a global scale. Journal of Hydrology 212–213: 268–286.

Ritchie (1972). Model for predicting evaporation from a row crop with incomplete cover. - Water resources research.

See Also

[spwb](#), [hydrology_soilWaterInputs](#)

Examples

```
SoilDepth = c(200,400,800,1200,1500)

#TOPSOIL LAYERS
d1 = pmin(SoilDepth, 300) #<300
#SUBSOIL LAYERS
d2 = pmax(0, pmin(SoilDepth-300,1200)) #300-1500 mm
#ROCK LAYER
d3 = 4000-(d1+d2) #From SoilDepth down to 4.0 m

TS_clay = 15
TS_sand = 25
SS_clay = 15
SS_sand = 25
RL_clay = 15
RL_sand = 25
TS_gravel = 20
```



```

SS_gravel = 40
RL_gravel = 95

Theta_FC1=soil_psi2thetaSX(TS_clay, TS_sand, -33) #in m3/m3
Theta_FC2=soil_psi2thetaSX(SS_clay, SS_sand, -33) #in m3/m3
Theta_FC3=soil_psi2thetaSX(RL_clay, RL_sand, -33) #in m3/m3
pcTS_gravel = 1-(TS_gravel/100)
pcSS_gravel = 1-(SS_gravel/100)
pcRL_gravel = 1-(RL_gravel/100)
MaxVol1 = (d1*Theta_FC1*pcTS_gravel)
MaxVol2 = (d2*Theta_FC2*pcSS_gravel)
MaxVol3 = (d3*Theta_FC3*pcRL_gravel)
V = MaxVol1+MaxVol2+MaxVol3

par(mar=c(5,5,1,1), mfrow=c(1,2))
NP = seq(0,60, by=1)
plot(NP,hydrology_infiltrationAmount(NP, V[1]), type="l", xlim=c(0,60), ylim=c(0,60),
      ylab="Infiltration (mm)", xlab="Net rainfall (mm)", frame=FALSE)
lines(NP,hydrology_infiltrationAmount(NP, V[2]), lty=2)
lines(NP,hydrology_infiltrationAmount(NP, V[3]), lty=3)
lines(NP,hydrology_infiltrationAmount(NP, V[4]), lty=4)
lines(NP,hydrology_infiltrationAmount(NP, V[5]), lty=5)
legend("topleft", bty="n", lty=1:5,
      legend=c(paste("d =", SoilDepth, "Vsoil =",round(V),"mm")))
plot(NP,NP-hydrology_infiltrationAmount(NP, V[1]), type="l", xlim=c(0,60), ylim=c(0,60),
      ylab="Runoff (mm)", xlab="Net rainfall (mm)", frame=FALSE)
lines(NP,NP-hydrology_infiltrationAmount(NP, V[2]), lty=2)
lines(NP,NP-hydrology_infiltrationAmount(NP, V[3]), lty=3)
lines(NP,NP-hydrology_infiltrationAmount(NP, V[4]), lty=4)
lines(NP,NP-hydrology_infiltrationAmount(NP, V[5]), lty=5)
legend("topleft", bty="n", lty=1:5,
      legend=c(paste("d =", SoilDepth,"Vsoil =",round(V),"mm")))

```

soil texture and hydraulics

Soil texture and hydraulics

Description

Collection of functions relating soil texture with soil hydraulics and soil water content (see description of each function in details).

Usage

```

soil_psi2thetaSX(clay, sand, psi, om = NA)
soil_psi2thetaVG(n, alpha, theta_res, theta_sat, psi)
soil_theta2psiSX(clay, sand, theta, om = NA)
soil_theta2psiVG(n, alpha, theta_res, theta_sat, theta)

```

```

soil_saturatedConductivitySX(clay, sand, om = NA, mmol = TRUE)
soil_USDAType(clay, sand)
soil_vanGenuchtenParamsCarsel(soilType)
soil_vanGenuchtenParamsToth(clay, sand, om, bd, topsoil)
soil_psi(soil, model="SX")
soil_theta(soil, model="SX")
soil_water(soil, model="SX")
soil_waterFC(soil, model="SX")
soil_waterWP(soil, model="SX")
soil_waterSAT(soil, model="SX")
soil_waterExtractable(soil, model="SX", minPsi = -5.0)
soil_thetaFC(soil, model="SX")
soil_thetaWP(soil, model="SX")
soil_thetaSAT(soil, model="SX")
soil_thetaSATSX(clay, sand, om = NA)
soil_waterTableDepth(soil, model="SX")
soil_rockWeight2Volume(pWeight, bulkDensity, rockDensity = 2.3)
soil_retentionCurvePlot(soil, model="SX", layer = 1,
                        psi = seq(0, -6.0, by=-0.01),
                        relative = TRUE, to = "SAT")

```

Arguments

clay	Percentage of clay (in percent weight).
sand	Percentage of sand (in percent weight).
n, alpha, theta_res, theta_sat	Parameters of the Van Genuchten-Mualem model ($m = 1 - 1/n$).
psi	Water potential (in MPa).
theta	Relative water content (in percent volume).
om	Percentage of organic matter (optional, in percent weight).
mmol	Boolean flag to indicate that saturated conductivity units should be returned in mmol/m/s/MPa. If mmol = FALSE then units are cm/day.
bd	Bulk density (in g/cm ³).
topsoil	A boolean flag to indicate topsoil layer.
soilType	A string indicating the soil type.
soil	Soil object (returned by function soil).
model	Either 'SX' or 'VG' for Saxton's or Van Genuchten's water retention models; or 'both' to plot both retention models.
layer	Soil layer to be plotted.
relative	Boolean flag to indicate that retention curve should be relative to field capacity or saturation.
to	Either 'SAT' (saturation) or 'FC' (field capacity).
minPsi	Minimum water potential (in MPa) to calculate the amount of extractable water.
pWeight	Percentage of corresponding to rocks, in weight.
bulkDensity	Bulk density of the soil fraction (g/cm ³).
rockDensity	Rock density (g/cm ³).

Details

- `soil_psi2thetaSX()` and `soil_theta2psiSX()` calculate water potentials (MPa) and water contents (theta) using texture data the formulae of Saxton et al. (1986) or Saxton & Rawls (2006) depending on whether organic matter is available.
- `soil_psi2thetaVG()` and `soil_theta2psiVG()` to the same calculations as before, but using the Van Genuchten - Mualem equations (Wösten & van Genuchten 1988).
- `soil_saturatedConductivitySX()` returns the saturated conductivity of the soil (in cm/day or mmol/m/s/MPa), estimated from formulae of Saxton et al. (1986) or Saxton & Rawls (2006) depending on whether organic matter is available.
- `soil_USDAType()` returns the USDA type (a string) for a given texture.
- `soil_vanGenuchtenParamsCarsel()` gives parameters for van Genuchten-Mualem equations (alpha, n, theta_res and theta_sat, where alpha is in MPa-1) for a given texture type (Leij et al. 1996)
- `soil_vanGenuchtenParamsToth()` gives parameters for van Genuchten-Mualem equations (alpha, n, theta_res and theta_sat, where alpha is in MPa-1) for a given texture, organic matter and bulk density (Toth et al. 2015).
- `soil_psi()` returns the water potential (MPa) of each soil layer, according to its water retention model.
- `soil_theta()` returns the moisture content (as percent of soil volume) of each soil layer, according to its water retention model.
- `soil_water()` returns the water volume (mm) of each soil layer, according to its water retention model.
- `soil_waterExtractable()` returns the water volume (mm) extractable from the soil according to its water retention curves and up to a given soil water potential.
- `soil_waterFC()` and `soil_thetaFC()` calculate the water volume (in mm) and moisture content (as percent of soil volume) of each soil layer at field capacity, respectively.
- `soil_waterWP()` and `soil_thetaWP()` calculate the water volume (in mm) and moisture content (as percent of soil volume) of each soil layer at wilting point (-1.5 MPa), respectively.
- `soil_waterSAT()`, `soil_thetaSATSX()` and `soil_thetaSAT()` calculate the saturated water volume (in mm) and moisture content (as percent of soil volume) of each soil layer.
- `soil_waterTableDepth()` returns water table depth in mm from surface.
- `soil_rockWeight2Volume()` transforms rock percentage from weight to volume basis.
- `soil_retentionCurvePlot()` allows plotting the water retention curve of a given soil layer.

Value

Depends on the function (see details).

Author(s)

Miquel De Cáceres Ainsa, CTFC

References

- Leij, F.J., Alves, W.J., Genuchten, M.T. Van, Williams, J.R., 1996. The UNSODA Unsaturated Soil Hydraulic Database User's Manual Version 1.0.
- Saxton, K.E., Rawls, W.J., Romberger, J.S., Papendick, R.I., 1986. Estimating generalized soil-water characteristics from texture. *Soil Sci. Soc. Am. J.* 50, 1031–1036.
- Saxton, K.E., Rawls, W.J., 2006. Soil water characteristic estimates by texture and organic matter for hydrologic solutions. *Soil Sci. Soc. Am. J.* 70, 1569. doi:10.2136/sssaj2005.0117
- Wösten, J.H.M., & van Genuchten, M.T. 1988. Using texture and other soil properties to predict the unsaturated soil hydraulic functions. *Soil Science Society of America Journal* 52: 1762–1770.
- Tóth, B., Weynants, M., Nemes, A., Makó, A., Bilas, G., & Tóth, G. 2015. New generation of hydraulic pedotransfer functions for Europe. *European Journal of Soil Science* 66: 226–238.

See Also

[soil](#)

Examples

```
#Determine USDA soil texture type
type = soil_USDAType(clay=40, sand=10)
type

#Van Genuchten's params (bulk density = 1.3 g/cm)
vg = soil_vanGenuchtenParamsToth(40,10,1,1.3,TRUE)
vg

# Initialize soil object with default params
s = soil(defaultSoilParams())

# Plot Saxton's and Van Genuchten's water retention curves
soil_retentionCurvePlot(s, model="both")
```

soil thermodynamics *Soil thermodynamic functions*

Description

Functions `soil_thermalConductivity` and `soil_thermalCapacity` calculate thermal conductivity and thermal capacity for each soil layer, given its texture and water content. Functions `soil_temperatureGradient` and `soil_temperatureChange` are used to calculate soil temperature gradients (in °C/m) and temporal temperature change (in °C/s) given soil layer texture and water content (and possibly including heat flux from above).

Usage

```

soil_thermalConductivity(soil, model = "SX")
soil_thermalCapacity(soil, model = "SX")
soil_temperatureChange(dVec, Temp, sand, clay, W, Theta_FC, Gdown)
soil_temperatureGradient(dVec, Temp)

```

Arguments

soil	Soil object (returned by function soil).
model	Either 'SX' or 'VG' for Saxton's or Van Genuchten's pedotransfer models.
dVec	Width of soil layers (in mm).
Temp	Temperature (in °C) for each soil layer.
clay	Percentage of clay (in percent weight) for each layer.
sand	Percentage of sand (in percent weight) for each layer.
W	Soil moisture (in percent of field capacity) for each layer.
Theta_FC	Relative water content (in percent volume) at field capacity for each layer.
Gdown	Downward heat flux from canopy to soil (in W·m ⁻²).

Value

Function `soil_thermalConductivity` returns a vector with values of thermal conductivity (W/m^{°K}) for each soil layer. Function `soil_thermalCapacity` returns a vector with values of heat storage capacity (J/m³/°K) for each soil layer. Function `soil_temperatureGradient` returns a vector with values of temperature gradient between consecutive soil layers. Function `soil_temperatureChange` returns a vector with values of instantaneous temperature change (°C/s) for each soil layer.

Author(s)

Miquel De Cáceres Ainsa, CTFC

References

- Cox, P.M., Betts, R.A., Bunton, C.B., Essery, R.L.H., Rowntree, P.R., & Smith, J. 1999. The impact of new land surface physics on the GCM simulation of climate and climate sensitivity. *Climate Dynamics* 15: 183–203.
- Dharssi, I., Vidale, P.L., Verhoef, A., MacPherson, B., Jones, C., & Best, M. 2009. New soil physical properties implemented in the Unified Model at PS18. 9–12.

See Also

[soil](#)

Examples

```

examplesoil = soil(defaultSoilParams())
soil_thermalConductivity(examplesoil)
soil_thermalCapacity(examplesoil)

#Values change when altering water content (drier layers have lower conductivity and capacity)
examplesoil$W = c(0.1, 0.4, 0.7, 1.0)
soil_thermalConductivity(examplesoil)
soil_thermalCapacity(examplesoil)

```

soilgridsParams	<i>SoilGrids soil description fetcher</i>
-----------------	---

Description

soilgridsParams takes a vector of depths and returns a list of soil characteristics ready to use with [soil](#) function.

Usage

```
soilgridsParams(points, widths = c(300, 700, 1000, 2000), verbose = FALSE)
```

Arguments

points	An object of class (or subclass) SpatialPoints with a valid CRS definition.
widths	A numeric vector indicating the desired layer widths, in <i>mm</i> . If NULL the default soil grids layer definition is returned.
verbose	A logical flag to include a progress bar while processing the output of the query to the SoilGrids REST API.

Details

This function connects with the SoilGrids REST API (<https://rest.soilgrids.org>) to retrieve the soil physical and chemical characteristics for a site (Hengl *et al.* 2007), selected by its coordinates. Also, in case the depths are not the default ones in the SoilGrids API, the function uses averages the values of soil grid layers depending on the overlap between soil layer definitions. Input coordinates are transformed to longitude/latitude within the function.

Value

If only one point is supplied, a data frame containing the soil characteristics ready to use with the [soil](#) function. If more than one point is supplied, the function returns a list with as many elements as points, each one containing the mentioned list.

Author(s)

Víctor Granda, CREA/CTFC
 Miquel De Cáceres Ainsa, CTFC

References

Hengl T, Mendes de Jesus J, Heuvelink GBM, Ruiperez Gonzalez M, Kilibarda M, Blagotić A, et al. (2017) SoilGrids250m: Global gridded soil information based on machine learning. PLoS ONE 12(2): e0169748. doi:10.1371/journal.pone.0169748.

See Also

[soil](#), [defaultSoilParams](#)

Examples

```
## Not run:
  coords_sp <- SpatialPoints(cbind(long = -5.6333, lat = 42.6667),
                             CRS("+proj=longlat +datum=WGS84"))
  foo <- soilgridsParams(coords_sp, widths = c(300, 700, 1000))
  foo_soil <- soil(foo)
  foo_soil

## End(Not run)
```

Species values

Species description functions

Description

Functions to calculate attributes of a [forest](#) object by species.

Usage

```
species_basalArea(x, SpParams)
species_cover(x, SpParams)
species_density(x, SpParams, mode = "MED")
species_foliarBiomass(x, SpParams, gdd = NA, mode = "MED")
species_fuel(x, SpParams, gdd = NA, includeDead = TRUE, mode = "MED")
species_LAI(x, SpParams, gdd = NA, mode = "MED")
species_phytovolume(x, SpParams)
```

Arguments

x	An object of class forest .
SpParams	A data frame with species parameters (see SpParamsMED).
gdd	Growth degree days (to account for leaf phenology effects).
includeDead	A flag to indicate that standing dead fuels (dead branches) are included.
mode	Calculation mode, either "MED" or "US".

Value

A vector with values for each species in SpParams:

- `species_basalArea`: Species basal area (m²/ha).
- `species_cover`: Shrub cover (in percent).
- `species_density`: Plant density (ind/ha). Tree density is directly taken from the forest object, while the shrub density is estimated from cover and height by calculating the area of a single individual.
- `species_foliarBiomass`: Standing biomass of leaves (in kg/m²).
- `species_fuel`: Fine fuel load (in kg/m²).
- `species_LAI`: Leaf area index (m²/m²).
- `species_phytovolume`: Shrub phytovolume (m³/m²).

Author(s)

Miquel De Cáceres Ainsa, CTFC

See Also

[spwb](#), [forest](#), [plant_basalArea](#), [summary.forest](#)

Examples

```
#Default species parameterization
data(SpParamsMED)

#Load example plot
data(exampleforestMED)

#Species basal area
species_basalArea(exampleforestMED, SpParamsMED)
```

SpParams

Data tables with species parameter values for Mediterranean and US

Description

A data sets of species parameter values, resulting from bibliographic search, fit to empirical data or expert-based guesses.

Usage

```
data("SpParamsMED")
data("SpParamsUS")
```


Format

A data frame with species as rows and the following columns (variables), grouped by categories.

Plant identity:

- Name: A factor with taxon names (mostly species names).
- IFNcodes: A factor with levels corresponding to species codes in the Third Spanish forest inventory (DGCN 2005).
- SpIndex: A numeric vector of the species index.
- Group: Either "Gymnosperm" or "Angiosperm".
- Order: Taxonomical order.
- Family: Taxonomical family.
- GrowthForm: Either Shrub, Tree and Tree/Shrub.

Plant size:

- Hmax: Maximum plant height (in cm).
- Hmed: Median plant height (in cm).
- Z50: Rooting depth (in mm) corresponding to 50% of fine roots.
- Z95: Rooting depth (in mm) corresponding to 95% of fine roots.

Shrub allometric coefficients (calibrated for Catalonia):

- a_ash: Regression coefficient relating the square of shrub height with shrub area.
- a_bsh, b_bsh: Allometric coefficients relating phytovolume with dry weight of shrub individuals.
- cr: Ratio between shrub crown length and total height.

Tree allometric coefficients (calibrated for Catalonia):

- a_fbt, b_fbt, c_fbt, d_fbt: Regression coefficients used to calculate foliar biomass of an individual tree from its dbh and the cumulative basal area of larger trees.
- a_cr, b_1cr, b_2cr, b_3cr, c_1cr, c_2cr: Regression coefficients used to calculate crown ratio of trees.
- a_cw, b_cw: Regression coefficients used to calculate crown width of trees.

Phenology:

- PhenologyType: Leaf phenology type, either "oneflush-evergreen" (new leaves develop in spring-summer), "progressive-evergreen" (new leaves develop during any season), "winter-deciduous" (leaf senescence in autumn, new leaves in spring-summer) or "winter-semideciduous" (same as before, but abscission of senescent leaves occurs when new leaves are produced).
- LeafDuration: Leaf duration (in years).
- Sgdd: Growth degree days for leaf budburst (in Celsius).

Anatomy:

- SLA: Specific leaf area ($\text{mm}^2/\text{mg} = \text{m}^2/\text{kg}$).
- LeafDensity: Density of leaf tissue (dry weight over volume).
- WoodDensity: Wood density (at 0 percent humidity!).
- FineRootDensity: Density of fine root tissue (dry weight over volume).
- r635: Ratio between the weight of leaves plus branches and the weight of leaves alone for branches of 6.35 mm.

- pDead: Proportion of total fine fuels that are dead
- A12As: Leaf area to sapwood area ratio (in $m^2 \cdot m^{-2}$).
- LeafWidth: Leaf width (in cm).
- SRL: Specific root length (cm/g).
- RLD: Fine root length density (i.e. density of root length per soil volume) ($cm \cdot cm^{-3}$).

Tissue moisture:

- maxFMC: Maximum fuel moisture (in percent of dry weight)
- minFMC: Minimum fuel moisture (in percent of dry weight)
- LeafPI0: Osmotic potential at full turgor of leaves (MPa).
- LeafEPS: Modulus of elasticity (capacity of the cell wall to resist changes in volume in response to changes in turgor) of leaves (MPa).
- LeafAF: Apoplastic fraction (proportion of water outside the living cells) in leaves.
- StemPI0: Osmotic potential at full turgor of symplastic xylem tissue (MPa).
- StemEPS: Modulus of elasticity (capacity of the cell wall to resist changes in volume in response to changes in turgor) of symplastic xylem tissue (Mpa).
- StemAF: Apoplastic fraction (proportion of water outside the living cells) in stem xylem.

Flammability:

- LigninPercent: Percent of lignin+cutin over dry weight in leaves.
- ParticleDensity: Particle density (kg/m^3).
- LeafLitterFuelType: Fuel type for leaf litter, with levels Broadleaved, LongLinear, Scale and ShortLinear.
- Flammability: Flammability modifier (either 1 or 2 for normal or high, respectively).
- SAV: Surface-area-to-volume ratio of the small fuel (1h) fraction (leaves and branches < 6.35mm) (m^2/m^3).
- HeatContent: High fuel heat content (kJ/kg).

Extinction:

- gammaSWR: Reflectance (albedo) for SWR ($\gamma_{PAR} = 0.8 \cdot \gamma_{SWR}$).
- alphaSWR: Absorbance coefficient for SWR ($\alpha_{PAR} = \alpha_{SWR} \cdot 1.35$).
- kPAR: Extinction coefficient for PAR ($k_{SWR} = k_{PAR} / 1.35$).
- g: Canopy water storage capacity per LAI unit (in mm/LAI).

Transpiration/Photosynthesis:

- WUE: Water use efficiency for carbon assimilation ($g \text{ C} / mm \text{ water}$).
- Psi_Extract: Water potential corresponding to 50% stomatal conductance (in MPa).
- Psi_Critic: Water potential corresponding to 50% of stem cavitation (in MPa).
- pRootDisc: Relative root conductance leading to hydraulic disconnection from a soil layer.
- Gwmin: Minimum stomatal conductance to water vapor per leaf area unit (in $mol \cdot s^{-1} \cdot m^{-2}$).
- Gwmax: Maximum stomatal conductance to water vapor per leaf area unit (in $mol \cdot s^{-1} \cdot m^{-2}$).
- VCleaf_kmax: Leaf hydraulic conductance (in $mmol \text{ H}_2\text{O} \cdot s^{-1} \cdot m^{-2} \cdot MPa^{-1}$).
- VCleaf_c, VCleaf_d: Parameters of the leaf vulnerability curve (VCleaf_d in MPa).

- `Kmax_stemxylem`: Sapwood-specific hydraulic conductivity of stem xylem (in $\text{kg H}_2\text{O}\cdot\text{s}^{-1}\cdot\text{m}^{-1}\cdot\text{MPa}^{-1}$).
- `VCstem_c`, `VCstem_d`: Parameters of the stem xylem vulnerability curve (`VCstem_d` in MPa).
- `Kmax_rootxylem`: Sapwood-specific hydraulic conductivity of root xylem (in $\text{kg H}_2\text{O}\cdot\text{s}^{-1}\cdot\text{m}^{-1}\cdot\text{MPa}^{-1}$).
- `VCroot_c`, `VCroot_d`: Parameters of the root xylem vulnerability curve (`VCroot_d` in MPa).
- `Narea`: Nitrogen concentration per leaf area (in $\text{g}\cdot\text{m}^{-2}$).
- `Vmax298`: Maximum Rubisco carboxylation rate at 25°C (in $\text{micromol CO}_2\cdot\text{s}^{-1}\cdot\text{m}^{-2}$).
- `Jmax298`: Maximum electron transport rate at 25°C (in $\text{micromol electrons}\cdot\text{s}^{-1}\cdot\text{m}^{-2}$).

Growth:

- `WoodC`: Wood carbon content per dry weight (g C / g dry).
- `RGRsapwoodmax`: Maximum relative growth rate (in basal area or sapwood area) (in $\text{cm}^2\cdot\text{cm}^{-2}$).
- `fHDmin`: Minimum value of the height-to-diameter ratio (dimensionless).
- `fHDmax`: Maximum value of the height-to-diameter ratio (dimensionless).

Details

See details of parameterization in De Caceres et al. (2015) and De Caceres et al. (submitted).

Source

De Cáceres M, Martínez-Vilalta J, Coll L, Llorens P, Casals P, Poyatos R, Pausas JG, Brotons L. (2015) Coupling a water balance model with forest inventory data to predict drought stress: the role of forest structural changes vs. climate changes. *Agricultural and Forest Meteorology* (doi:10.1016/j.agrformet.2015.06.012).

DGCN (2005). Tercer Inventario Forestal Nacional (1997-2007): Catalunya. Dirección General de Conservación de la Naturaleza, Ministerio de Medio Ambiente, Madrid.

See Also

[spwb](#)

Examples

```
data(SpParamsMED)
```

spwb

*Soil-plant water balance***Description**

Function `spwb()` is a water balance model that determines changes in soil moisture, soil water potentials, plant transpiration and drought stress at daily steps for a given forest stand during a period specified in the input climatic data. Function `pwb()` performs plant water balance only (i.e. soil moisture dynamics is an input) at daily steps for a given forest stand during a period specified in the input climatic data. On both simulation functions plant transpiration and photosynthesis processes are conducted with different level of detail depending on the transpiration mode.

Usage

```
spwb(x, soil, meteo, latitude, elevation = NA, slope = NA, aspect = NA)
pwb(x, soil, meteo, W, latitude, elevation = NA, slope = NA, aspect = NA,
    canopyEvaporation = numeric(0),
    snowMelt = numeric(0),
    soilEvaporation = numeric(0))
```

Arguments

- | | |
|-------|--|
| x | An object of class <code>spwbInput</code> . |
| soil | A list containing the description of the soil (see <code>soil</code>). |
| meteo | <p>A data frame with daily meteorological data series. Row names of the data frame should correspond to date strings with format "yyyy-mm-dd" (see <code>Date</code>). When using the 'Granier' transpiration mode the following columns are required:</p> <ul style="list-style-type: none"> • MeanTemperature: Mean temperature (in degrees Celsius). • Precipitation: Precipitation (in mm). • Radiation: Solar radiation (in MJ/m2/day), required only if <code>snowpack = TRUE</code>. • PET: Potential evapotranspiration (in mm). <p>When using the 'Sperry' transpiration mode the following columns are required:</p> <ul style="list-style-type: none"> • MeanTemperature: Mean temperature (in degrees Celsius). • MinTemperature: Minimum temperature (in degrees Celsius). • MaxTemperature: Maximum temperature (in degrees Celsius). • MinRelativeHumidity: Minimum relative humidity (in percent). • MaxRelativeHumidity: Maximum relative humidity (in percent). • Precipitation: Precipitation (in mm). • Radiation: Solar radiation (in MJ/m2/day). • WindSpeed: Wind speed (in m/s). If not available, this column can be left with NA values. |
| W | A matrix with the same number of rows as <code>meteo</code> and as many columns as soil layers, containing the soil moisture of each layer as proportion of field capacity. |

latitude	Latitude (in degrees).
elevation, slope, aspect	Elevation above sea level (in m), slope (in degrees) and aspect (in degrees from North). Required when using the 'Sperry' transpiration mode. Elevation is also required for 'Granier' if snowpack dynamics are simulated.
canopyEvaporation	A vector of daily canopy evaporation (from interception) values (mm). The length should match the number of rows in meteo.
soilEvaporation	A vector of daily bare soil evaporation values (mm). The length should match the number of rows in meteo.
snowMelt	A vector of daily snow melt values (mm). The length should match the number of rows in meteo.

Details

The model using 'Granier' transpiration mode is illustrated by function [transp_transpirationGranier](#) and described in De Caceres et al. (2015). Simulations using the 'Sperry' transpiration mode are computationally much more expensive, are described in De Cáceres et al. (2021) and are illustrated by function [transp_transpirationSperry](#).

Value

Function `spwb` returns a list of class 'spwb' whereas Function `pwb` returns a list of class 'pwb'. There are many elements in common in these lists, so they are listed here together:

- "latitude": Latitude (in degrees) given as input.
- "topography": Vector with elevation, slope and aspect given as input.
- "spwbInput": An (unmodified) copy of the object `x` of class [spwbInput](#) given as input (note that `x` is modified by the simulation function).
- "soilInput": An (unmodified) copy of the object `soil` of class [soil](#) given as input (note that `soil` is modified by the simulation function).
- "WaterBalance": A data frame where different variables (in columns) are given for each simulated day (in rows):
 - "PET": Potential evapotranspiration (in mm).
 - "Precipitation": Input precipitation (in mm).
 - "Rain": Precipitation as rain (in mm).
 - "Snow": Precipitation as snow (in mm).
 - "NetRain": Net rain, after accounting for interception (in mm).
 - "Infiltration": The amount of water infiltrating into the soil (in mm).
 - "Runoff": The amount of water exported via surface runoff (in mm).
 - "DeepDrainage": The amount of water exported via deep drainage (in mm).
 - "Evapotranspiration": Evapotranspiration (in mm).
 - "SoilEvaporation": Bare soil evaporation (in mm).

- "PlantExtraction": Amount of water extracted from soil by plants (in mm) (can only be different from transpiration for `transpirationMode = "Sperry"` when capacitance is considered).
- "Transpiration": Plant transpiration (considering all soil layers) (in mm).
- "HydraulicRedistribution": Water redistributed among soil layers, transported through the plant hydraulic network (only for `transpirationMode = "Sperry"`).
- "EnergyBalance": A data frame with the daily values of energy balance components for the soil and the canopy (only for `transpirationMode = "Sperry"`).
- "Temperature": A data frame with the daily values of minimum/mean/maximum temperatures for the atmosphere (input), canopy and soil (only for `transpirationMode = "Sperry"`).
- "Soil": A data frame where different variables (in columns) are given for each simulated day (in rows):
 - "W.1", ..., "W.k": Relative soil moisture content (relative to field capacity) in each soil layer.
 - "ML.1", ..., "ML.k": Soil water volume in each soil layer (in L/m²).
 - "MLTot": Total soil water volume (in L/m²).
 - "SWE": Snow water equivalent (mm) of the snow pack.
 - "PlantExt.1", ..., "PlantExt.k": Plant extraction from each soil layer (in mm).
 - "HydraulicInput.1", ..., "HydraulicInput.k": Water that entered the layer coming from other layers and transported via the plant hydraulic network (in mm) (only for `transpirationMode = "Sperry"`).
 - "psi.1", ..., "psi.k": Soil water potential in each soil layer (in MPa).
- "Stand": A data frame where different variables (in columns) are given for each simulated day (in rows):
 - "LAI": LAI of the stand (including live and dead leaves) (in m²/m²).
 - "LAIlive": LAI of the stand assuming all leaves are unfolded (in m²/m²).
 - "LAIexpanded": LAI of the stand of leaves actually unfolded (in m²/m²).
 - "LAIdead": LAI of the stand corresponding to dead leaves (in m²/m²).
 - "Cm": Water retention capacity of the canopy (in mm) (accounting for leaf phenology).
 - "LgroundPAR": The proportion of PAR that reaches the ground (accounting for leaf phenology).
 - "LgroundSWR": The proportion of SWR that reaches the ground (accounting for leaf phenology).
- "Plants": A list of daily results for plant cohorts (see below).
- "subdaily": A list of objects of class `spwb_day`, one per day simulated (only if required in control parameters, see `defaultControl`).

When `transpirationMode = "Granier"`, element "Plants" is a list with the following subelements:

- "LAI": A data frame with the daily leaf area index for each plant cohort.
- "AbsorbedSWRFraction": A data frame with the fraction of SWR absorbed by each plant cohort.

- "Transpiration": A data frame with the amount of daily transpiration (in mm) for each plant cohort.
- "Photosynthesis": A data frame with the amount of daily net photosynthesis (in g C·m⁻²) for each plant cohort.
- "PlantPsi": A data frame with the average daily water potential of each plant (in MPa).
- "PlantStress": A data frame with the amount of daily stress [0-1] suffered by each plant cohort (relative whole-plant conductance).

If transpirationMode="Sperry", element "Plants" is a list with the following subelements:

- "LAI": A data frame with the daily leaf area index for each plant cohort.
- "AbsorbedSWR": A data frame with the daily SWR absorbed by each plant cohort.
- "AbsorbedLWR": A data frame with the daily LWR absorbed by each plant cohort.
- "Transpiration": A data frame with the amount of daily transpiration (in mm) for each plant cohorts.
- "GrossPhotosynthesis": A data frame with the amount of daily gross photosynthesis (in g C·m⁻²) for each plant cohort.
- "NetPhotosynthesis": A data frame with the amount of daily net photosynthesis (in g C·m⁻²) for each plant cohort.
- "dEdP": A data frame with mean daily values of soil-plant conductance (derivative of the supply function) for each plant cohort.
- "PlantWaterBalance": A data frame with the daily balance between transpiration and soil water extraction for each plant cohort.
- "SunlitLeaves" and "ShadeLeaves": A list with daily results for sunlit and shade leaves:
 - "PsiMin": A data frame with the minimum (midday) daily sunlit or shade leaf water potential (in MPa).
 - "PsiMax": A data frame with the maximum (predawn) daily sunlit or shade leaf water potential (in MPa).
- "LeafPsiMin": A data frame with the minimum (midday) daily (average) leaf water potential of each plant (in MPa).
- "LeafPsiMax": A data frame with the maximum (predawn) daily (average) leaf water potential of each plant (in MPa).
- "LeafRWC": A data frame with the average daily leaf relative water content of each plant (in percent).
- "StemRWC": A data frame with the average daily stem relative water content of each plant (in percent).
- "LeafSympRWC": A data frame with the average daily leaf symplasm relative water content of each plant (in percent).
- "StemSympRWC": A data frame with the average daily stem symplasm relative water content of each plant (in percent).
- "StemPsi": A data frame with the minimum daily stem water potential of each plant (in MPa).
- "StemPLC": A data frame with the average daily proportion of stem conductance loss of each plant ([0-1]).

- "RootPsi": A data frame with the minimum daily root water potential of each plant (in MPa).
- "RhizoPsi": A list of data frames (one per plant cohort) with the minimum daily root water potential of each plant (in MPa).
- "PlantStress": A data frame with the amount of daily stress [0-1] suffered by each plant cohort (relative whole-plant conductance).

Note

State variables stored in objects `x` and `soil` are modified during the simulation.

Author(s)

Miquel De Cáceres Ainsa, CTFC

References

De Cáceres M, Martínez-Vilalta J, Coll L, Llorens P, Casals P, Poyatos R, Pausas JG, Brotons L. (2015) Coupling a water balance model with forest inventory data to predict drought stress: the role of forest structural changes vs. climate changes. *Agricultural and Forest Meteorology* 213: 77-90 (doi:10.1016/j.agrformet.2015.06.012).

De Cáceres M, Mencuccini M, Martin-StPaul N, Limousin JM, Coll L, Poyatos R, Cabon A, Granda V, Forner A, Valladares F, Martínez-Vilalta J (2021) Unravelling the effect of species mixing on water use and drought stress in holm oak forests: a modelling approach. *Agricultural and Forest Meteorology* 296 (doi:10.1016/j.agrformet.2020.108233).

See Also

[spwbInput](#), [spwb_day](#), [plot.spwb](#), [spwb_ldrOptimization](#), [forest](#)

Examples

```
#Load example daily meteorological data
data(examplemeteo)

#Load example plot plant data
data(exampleforestMED)

#Default species parameterization
data(SpParamsMED)

#Initialize soil with default soil params (4 layers)
examplesoil1 = soil(defaultSoilParams(4))

#Initialize control parameters
control = defaultControl()

#Initialize input
x1 = forest2spwbInput(exampleforestMED,examplesoil1, SpParamsMED, control)

#Call simulation function
```



```

S1<-spwb(x1, examplesoil1, examplometeo, latitude = 41.82592, elevation = 100)

#Plot results
plot(S1)

#Monthly summary (averages) of soil water balance
summary(S1, freq="months",FUN=mean, output="Soil")

## Not run:
examplesoil2 = soil(defaultSoilParams(4))

#Switch to 'Sperry' transpiration mode
control$transpirationMode="Sperry"

#Initialize input
x2 = forest2spwbInput(exampleforestMED,examplesoil2, SpParamsMED, control)

#Call simulation function (11 days)
d = 100:110
S2<-spwb(x2, examplesoil2, examplometeo[d,], latitude = 41.82592, elevation = 100)

# Run the model with 'Sperry' transpiration mode using the water balance of
# simulated with the 'Granier' model
P2<-pwb(x2, examplesoil2, examplometeo[d,], latitude = 41.82592, elevation = 100,
        W = as.matrix(S1$Soil[d, c("W.1", "W.2")]),
        canopyEvaporation = S1$WaterBalance$Interception[d],
        snowMelt = S1$WaterBalance$Snowmelt[d],
        soilEvaporation = S1$WaterBalance$SoilEvaporation[d])

## End(Not run)

```

spwb_day

Single day simulation

Description

Function `spwb_day` performs water balance for a single day and `growth_day` performs water and carbon balance for a single day.

Usage

```

spwb_day(x, soil, date, tmin, tmax, rhmin, rhmax, rad, wind,
         latitude, elevation, slope, aspect, prec, runon = 0.0)
growth_day(x, soil, date, tmin, tmax, rhmin, rhmax, rad, wind,
          latitude, elevation, slope, aspect, prec, runon = 0.0)

```

Arguments

x	An object of class spwbInput or growthInput .
soil	A list containing the description of the soil (see soil).
date	Date as string "yyyy-mm-dd".
tmin, tmax	Minimum and maximum temperature (in degrees Celsius).
rhmin, rhmax	Minimum and maximum relative humidity (in percent).
rad	Solar radiation (in MJ/m2/day).
wind	Wind speed (in m/s).
prec	Precipitation (in mm).
latitude	Latitude (in degrees). Required when using the 'Sperry' transpiration mode.
elevation, slope, aspect	Elevation above sea level (in m), slope (in degrees) and aspect (in degrees from North). Required when using the 'Sperry' transpiration mode.
runon	Surface water amount running on the target area from upslope (in mm).

Details

Detailed model description is available in the vignettes section. The model using 'Granier' transpiration mode is described in De Caceres et al. (2015). Simulations using the 'Sperry' transpiration mode are computationally much more expensive, are described in De Cáceres et al. (2021) and are illustrated by function [transp_transpirationSperry](#).

Value

Function `spwb_day()` returns a list of class `spwb_day` with the following elements:

- "cohorts": A data frame with cohort information, copied from [spwbInput](#).
- "WaterBalance": A vector of water balance components (rain, snow, net rain, infiltration, ...) for the simulated day, equivalent to one row of 'WaterBalance' object given in [spwb](#).
- "Soil": A data frame with results for each soil layer:
 - "SoilEvaporation": Water evaporated from the soil surface (in mm).
 - "HydraulicInput": Water entering each soil layer from other layers, transported via plant hydraulic network (in mm) (only for `transpirationMode = "Sperry"`).
 - "HydraulicOutput": Water leaving each soil layer (going to other layers or the transpiration stream) (in mm) (only for `transpirationMode = "Sperry"`).
 - "PlantExtraction": Water extracted by plants from each soil layer (in mm).
 - "psi": Soil water potential (in MPa).
- "Stand": A named vector with with stand values for the simulated day, equivalent to one row of 'Stand' object returned by [spwb](#).
- "Plants": A data frame of results for each plant cohort (see [transp_transpirationGranier](#) or [transp_transpirationSperry](#)).

The following items are only returned when `transpirationMode = "Sperry"`:

- "EnergyBalance": Energy balance of the stand (see [transp_transpirationSperry](#)).
- "RhizoPsi": Minimum water potential (in MPa) inside roots, after crossing rhizosphere, per cohort and soil layer.
- "SunlitLeaves" and "ShadeLeaves": For each leaf type, a data frame with values of LAI, Vmax298 and Jmax298 for leaves of this type in each plant cohort.
- "ExtractionInst": Water extracted by each plant cohort during each time step.
- "PlantsInst": A list with instantaneous (per time step) results for each plant cohort (see [transp_transpirationSperry](#)).
- "LightExtinction": A list of information regarding radiation balance through the canopy, as returned by function [light_instantaneousLightExtinctionAbsortion](#).
- "WindExtinction": A numeric vector with the wind speed (in m/s) for each plant cohort.

Note

Objects x and soil are modified during the simulation.

Author(s)

Miquel De Cáceres Ainsa, CTFC

References

De Cáceres M, Martínez-Vilalta J, Coll L, Llorens P, Casals P, Poyatos R, Pausas JG, Brotons L. (2015) Coupling a water balance model with forest inventory data to predict drought stress: the role of forest structural changes vs. climate changes. *Agricultural and Forest Meteorology* (doi:10.1016/j.agrformet.2015.06.012).

De Cáceres M, Mencuccini M, Martin-StPaul N, Limousin JM, Coll L, Poyatos R, Cabon A, Granda V, Forner A, Valladares F, Martínez-Vilalta J (2021) Unravelling the effect of species mixing on water use and drought stress in holm oak forests: a modelling approach. *Agricultural and Forest Meteorology* 296 (doi: 10.1016/j.agrformet.2020.108233).

See Also

[spwbInput](#), [spwb](#), [plot.spwb_day](#), [growthInput](#), [growth](#), [plot.growth_day](#)

Examples

```
#Load example daily meteorological data
data(examplemeteo)

#Load example plot plant data
data(exampleforestMED)

#Default species parameterization
data(SpParamsMED)

#Initialize control parameters
control = defaultControl()
```

```

# Day to be simulated
d = 100

#Simulate water balance one day only (Granier)
examplesoil = soil(defaultSoilParams(2))
x1 = forest2spwbInput(exampleforestMED,examplesoil, SpParamsMED, control)
sd1<-spwb_day(x1, examplesoil, rownames(examplemeteo)[d],
              examplemeteo$MinTemperature[d], examplemeteo$MaxTemperature[d],
              examplemeteo$MinRelativeHumidity[d], examplemeteo$MaxRelativeHumidity[d],
              examplemeteo$Radiation[d], examplemeteo$WindSpeed[d],
              latitude = 41.82592, elevation = 100, slope=0, aspect=0,
              prec = examplemeteo$Precipitation[d])

#Simulate water balance for one day only (Sperry's mode)
control$transpirationMode="Sperry"
examplesoil2 = soil(defaultSoilParams(2))
x2 = forest2spwbInput(exampleforestMED,examplesoil2, SpParamsMED, control)
sd2<-spwb_day(x2, examplesoil2, rownames(examplemeteo)[d],
              examplemeteo$MinTemperature[d], examplemeteo$MaxTemperature[d],
              examplemeteo$MinRelativeHumidity[d], examplemeteo$MaxRelativeHumidity[d],
              examplemeteo$Radiation[d], examplemeteo$WindSpeed[d],
              latitude = 41.82592, elevation = 100, slope=0, aspect=0,
              prec = examplemeteo$Precipitation[d])

#Plot plant transpiration (see function 'plot.swb.day()')
plot(sd2)

#Simulate water and carbon balance for one day only
examplesoil3 = soil(defaultSoilParams(2))
x3 = forest2growthInput(exampleforestMED,examplesoil3, SpParamsMED, control)
sd3<-growth_day(x3, examplesoil3, rownames(examplemeteo)[d],
                examplemeteo$MinTemperature[d], examplemeteo$MaxTemperature[d],
                examplemeteo$MinRelativeHumidity[d], examplemeteo$MaxRelativeHumidity[d],
                examplemeteo$Radiation[d], examplemeteo$WindSpeed[d],
                latitude = 41.82592, elevation = 100, slope=0, aspect=0,
                prec = examplemeteo$Precipitation[d])

```

spwb_ldrCalibration *Calibration of root distribution*

Description

The function `spwb_ldrCalibration` calibrates the species root distribution within `spwb`, given the arguments `x`, `meteo`, `soil`, `psi_crit`, `obs` and `calibVar`. This calibration is based on reference measured values. These reference measured values can be Soil water content, Total

transpiration or Transpiration by cohort. Return the calibrated root distribution for each tree species (no shrub calibration is done), expressed as parameters of the function `root_ldrDistribution`.

Usage

```
spwb_ldrCalibration(x, soil, meteo, calibVar, obs,
                   RZmin = 301, RZmax = 4000,
                   V1min = 0.01, V1max = 0.94, resolution = 20, heat_stop = 0,
                   transformation = "identity", verbose = FALSE)
```

Arguments

x	An object of class <code>spwbInput</code> .
soil	A list containing the description of the soil (see <code>soil</code>).
meteo	A data frame with daily meteorological data series. When using the 'Granier' transpiration mode the following columns are required: <ul style="list-style-type: none"> • DOY: Day of the year (Julian day). • Precipitation: Precipitation (in mm). • MeanTemperature: Mean temperature (in degrees Celsius). • PET: Potential evapotranspiration (in mm).
calibVar	A character string indicating the calibration variable to be used. It can be one of the following: SWC, Eplanttot or Cohorts.
obs	Measured calibration variable. Depending on the value of calibVar it can be a numeric vector with the measured SWC values (if calibVar = "SWC"), or a data frame with the first column containing the measured total transpiration (named Eplanttot) and the following columns containing the cohorts transpiration.
RZmin	The minimum value of RZ (the rooting depth) to be explored (in mm)
RZmax	The maximum value of RZ (the rooting depth) to be explored (in mm)
V1min	The minimum value of V1 (the root proportion in the first soil layer) to be explored
V1max	The maximum value of V1 (the root proportion in the first soil layer) to be explored
resolution	An integer defining the number of values to obtain by discretization of the root parameters RZ and V1. The number of parameter combinations and therefore the computation cost increases increase with the square of resolution
transformation	Function to modify the size of Z intervals to be explored (by default, bins are equal).
heat_stop	An integer defining the number of days during to discard from the calculation of the optimal root distribution. Usefull if the soil water content initialization is not certain
verbose	A logical value. Print the internal messages of the function?

Details

This function performs three different kinds of calibration, selecting those root distribution parameters that minimize the MAE between the predicted values and the measured values provided in `obs` argument. If `calibVar = "SWC"` different V1 values are tested running `spwb` maintaining the total soil depth provided in `x` and assuming that value is also the depth containing 95 percent of the roots. If `calibVar = "Eplanttot"` or `calibVar = 'Cohorts'` different combinations of RZ and V1 values are tested for each tree cohort and the root parameters are selected based on the MAE between the total transpiration or the cohort transpiration.

Value

The function returns a data frame containing the species index used in `medfate`, calibrated values for Z50, Z95 and V1 and the MAE value for that combination.

Author(s)

Víctor Granda, CTFC-CREAF

Antoine Cabon, CTFC

Miquel De Cáceres Ainsa, CTFC

See Also

[spwb_ldrOptimization](#) for when no measured data is available, [spwb](#), [soil](#), [root_ldrDistribution](#)

spwb_ldrOptimization *Optimization of root distribution*

Description

Functions `spwb_ldrExploration` and `spwb_ldrOptimization` are used to find optimum the species root distribution within `spwb`, given the arguments `x`, `meteo`, `soil` and `psi_crit`.

Usage

```
spwb_ldrExploration(x, soil, meteo, cohorts = NULL,
                    RZmin = 301, RZmax = 4000,
                    V1min = 0.01, V1max = 0.94, resolution = 10, heat_stop = 0,
                    transformation = "identity",
                    verbose = FALSE, ...)
spwb_ldrOptimization(y, psi_crit, opt_mode = 1)
```

Arguments

x	An object of class <code>spwbInput</code> .
soil	A list containing the description of the soil (see <code>soil</code>).
meteo	A data frame with daily meteorological data series (see <code>spwb</code>).
cohorts	A character string with the names of cohorts to be explored. If NULL then all cohorts are explored.
RZmin	The minimum value of RZ (the rooting depth) to be explored (in mm)
RZmax	The maximum value of RZ (the rooting depth) to be explored (in mm)
V1min	The minimum value of V1 (the root proportion in the first soil layer) to be explored
V1max	The maximum value of V1 (the root proportion in the first soil layer) to be explored
resolution	An integer defining the number of values to obtain by discretization of the root parameters RZ and V1. The number of parameter combinations and therefore the computation cost increases increase with the square of resolution
transformation	Function to modify the size of Z intervals to be explored (by default, bins are equal).
heat_stop	An integer defining the number of days during to discard from the calculation of the optimal root distribution. Usefull if the soil water content initialization is not certain
...	Additional parameters to function <code>spwb</code> .
y	The result of calling <code>spwb_ldrExploration</code> .
psi_crit	A numerical vector of length equal to the number of species in the plot containing the species values of water potential inducing hydraulic failure (in MPa). Use NA values to skip optimization for particular plant cohorts.
opt_mode	Optimization mode: <ul style="list-style-type: none"> • <code>opt_mode = 1</code> maximizes transpiration along the line of stress equal to <code>psi_crit</code> (Cabon et al. 2018). The optimization is based on the eco-hydrological equilibrium hypothesis (Eagleson, 1982), which is formulated here as the root distribution for which plant transpiration is maximized while the plant water potential is close to the species-defined critical value <code>psi_crit</code> (Cabon et al.,2018). • <code>opt_mode = 2</code> maximizes transpiration among combinations with stress according to <code>psi_crit</code>). • <code>opt_mode = 3</code> maximizes photosynthesis among combinations with stress according to <code>psi_crit</code>). • <code>opt_mode = 4</code> maximizes transpiration, subject to root construction constrains, among combinations with stress according to <code>psi_crit</code>). • <code>opt_mode = 5</code> maximizes photosynthesis, subject to root construction constrains, among combinations with stress according to <code>psi_crit</code>).
verbose	A logical value. Print the internal messages of the function?

Details

For each combination of the parameters RZ and V1 the function `spwb_ldrExploration` runs `spwb`, setting the total soil depth equal to RZ. The root proportion in each soil layer is derived from V1, the depth of the first soil layer (as defined in the argument `soil`) and RZ using the LDR root distribution model (Schenk and Jackson, 2002) and assuming that the depth containing 95 percent of the roots is equal to RZ. Function `spwb_ldrOptimization` takes the result of the exploration and tries to find optimum root distribution parameters. `psi_crit`, the species specific water potential inducing hydraulic failure, can be approached by the water potential inducing 50 percent of loss of conductance for the angiosperms and 88 percent for the gymnosperms (Urli et al., 2013, Brodrribb et al., 2010). Details of the hypothesis and limitations of the optimization method are given in Cabon et al. (2019).

Value

Function `spwb_ldrExploration` returns a list containing a list containing the explored RZ and V1 combinations as well as arrays with the values of average daily plant transpiration, average daily net photosynthesis and the minimum plant water potential for each cohort and parameter combination.

Function `spwb_ldrOptimization` returns a data frame with containing the species index used in `medfate`, `psi_crit` and the optimized values of V1 and the LDR parameters Z50 and Z95 (see [root_ldrDistribution](#)) and as many rows as the number of species.

Author(s)

Antoine Cabon, CTFC

Miquel De Cáceres Ainsa, CTFC

References

- Brodrribb, T.J., Bowman, D.J.M.S., Nichols, S., Delzon, S., Burlett, R., 2010. Xylem function and growth rate interact to determine recovery rates after exposure to extreme water deficit. *New Phytol.* 188, 533–542. doi:10.1111/j.1469-8137.2010.03393.x
- Cabon, A., Martínez-Vilalta, J., Poyatos, R., Martínez de Aragón, J., De Cáceres, M. (2018) Applying the eco-hydrological equilibrium hypothesis to estimate root distribution in water-limited forests. *Ecohydrology* 11: e2015.
- Eagleson, P.S., 1982. Ecological optimality in water-limited natural soil-vegetation systems: 1. Theory and hypothesis. *Water Resour. Res.* 18, 325–340. doi:10.1029/WR018i002p00325
- Schenk, H.J., Jackson, R.B., 2002. The Global Biogeography of Roots. *Ecol. Monogr.* 72, 311. doi:10.2307/3100092
- Urli, M., Porte, A.J., Cochard, H., Guengant, Y., Burlett, R., Delzon, S., 2013. Xylem embolism threshold for catastrophic hydraulic failure in angiosperm trees. *Tree Physiol.* 33, 672–683. doi:10.1093/treephys/tpt030

See Also

[spwb](#), [soil](#), [root_ldrDistribution](#)

Examples

```

## Not run:
#Load example daily meteorological data
data(examplemeteo)

#Load example plot plant data
data(exampleforestMED)

#Default species parameterization
data(SpParamsMED)

#Initialize soil with default soil params
examplesoil = soil(defaultSoilParams(2))

#Initialize control parameters
control = defaultControl()

#Initialize input
x = forest2spwbInput(exampleforestMED,examplesoil, SpParamsMED, control)

#Run exploration
y = spwb_ldrExploration(x = x, soil = examplesoil, meteo = examplemeteo,
                       elevation = 100)

#Optimization under different modes
spwb_ldrOptimization(y = y, psi_crit = c(-2,-3,-4), opt_mode = 1)
spwb_ldrOptimization(y = y, psi_crit = c(-2,-3,-4), opt_mode = 2)
spwb_ldrOptimization(y = y, psi_crit = c(-2,-3,-4), opt_mode = 3)

## End(Not run)

```

```

spwb_maximumTranspirationRatioPlot
      Maximum transpiration vs. LAI

```

Description

Builds a curve of maximum transpiration (Tmax) over potential evapotranspiration (ETP) for increasing leaf area index (LAI) values.

Usage

```

spwb_maximumTranspirationRatioPlot(x, soil, meteo, latitude, elevation, slope, aspect,
                                   ndays = 100,
                                   LAI_seq = c(0.1, 0.25, seq(0.5, 10, by = 0.5)))

```

Arguments

x	An object of class spwbInput .
soil	A list containing the description of the soil (see soil).
meteo	A data frame with daily meteorological data series.
latitude	Latitude (in degrees). Required when using the 'Sperry' transpiration mode.
elevation, slope, aspect	Elevation above sea level (in m), slope (in degrees) and aspect (in degrees from North). Required when using the 'Sperry' transpiration mode.
ndays	Number of days to subset for curve building.
LAI_seq	Sequence of stand LAI values to be tested.

Details

TO BE DONE

Author(s)

Miquel De Cáceres Ainsa, CTFC

See Also

[spwb](#)

Examples

```
## Not run:
#Load example daily meteorological data
data(examplemeteo)

#Load example plot plant data
data(exampleforestMED)

#Default species parameterization
data(SpParamsMED)

#Initialize soil with default soil params
examplesoil2 = soil(defaultSoilParams(2))

#Initialize control parameters
control = defaultControl()

#Switch to 'Sperry' transpiration mode
control$transpirationMode="Sperry"

#Initialize input
x2 = forest2spwbInput(exampleforestMED,examplesoil2, SpParamsMED, control)

#Run optimization
```

```

spwb_maximumTranspirationRatioPlot(x2, examplesoil2, examplemeteo,
                                   41.82592, elevation = 100,
                                   slope = 0, aspect = 0, ndays = 20)

## End(Not run)

```

spwb_resistances *Soil-plant resistances*

Description

Calculates and draws rhizosphere, root, stem and leaf resistances for simulation time steps

Usage

```

spwb_resistances(x, cohort = 1, relative = FALSE, draw = FALSE,
                 cumulative = FALSE, xlab = NULL, ylab = NULL)

```

Arguments

x	An object of class spwb. The function only works with the result of simulations with <code>transpirationMode = "Sperry"</code> .
cohort	An integer index indicating the cohort for which resistances are desired (by default the first cohort).
relative	A boolean flag to indicate that relative percentages are desired as output
draw	A boolean flag to indicate that a plot should be drawn.
cumulative	A flag to indicate that drawn series should be cumulative.
xlab	x-axis label.
ylab	y-axis label.

Details

The function makes internal calls to [hydraulics_soilPlantResistances](#).

Value

A data frame with dates in rows and resistance segments in columns (Rhizosphere, Root, Stem and Leaf). Values depend on whether `relative = TRUE` (percentages) or `relative = FALSE` (absolute resistance values). If `draw = TRUE` then a plot object is returned.

Author(s)

Miquel De Cáceres Ainsa, CTFC

See Also

[spwb](#), [plot.spwb](#)

spwb_sensitivity *Sensitivity analysis for soil plant water balance simulations*

Description

Performs a set of calls to [spwb](#) with the aim to determine the sensitivity to particular parameters.

Usage

```
spwb_sensitivity(x, soil, meteo,
                paramType = "above", paramName = "LAI_live", cohort = 1,
                p_change = c(-80,-40,-20,0,20,40,80),
                summary.fun = NULL, simplify=TRUE,...)
```

Arguments

x	An object of class spwbInput .
soil	A list containing the description of the soil (see soil).
meteo	A data frame with daily meteorological data series (see spwb).
paramType	Data frame of x to modify.
paramName	Name of the parameter to modify.
cohort	Integer with the cohort to modify.
p_change	Numerical vector with percentages of change.
summary.fun	Summary function to be applied to the results of each simulation.
simplify	Whether the result of summary.fun should be simplified (see sapply).
...	Additional parameters to function spwb .

Details

Due to parameter dependence, modifying some parameters affects others:

- Setting paramName = "Z50/Z95" affects below\$V, below\$VCroot_kmax and below\$VGrhizo_kmax.
- Modifying LAI_live also affects LAI_expanded.
- Modifying VCroot_kmax from paramsTranspiration affects both VCroot_kmax and below\$VCroot_kmax.
- Modifying WaterStorage affects simultaneously Vleaf and Vsapwood from paramsWaterStorage.
- Modifying c from paramsTranspiration affects simultaneously VCleaf_c, VCstem_c and VCroot_c.
- Modifying d from paramsTranspiration affects simultaneously VCleaf_d, VCstem_d and VCroot_d.
- Modifying Plant_kmax from paramsTranspiration affects VCleaf_kmax, VCstem_kmax, VCroot_kmax and below\$VCroot_kmax.
- Modifying Al2As from paramsAnatomy affects Vsapwood in paramsWaterStorage, VCstem_kmax and VCroot_kmax of paramsTranspiration and below\$VCroot_kmax.
- Setting paramName = "Vmax298/Jmax298" affects both Vmax298 and Jmax298 from paramsTranspiration.

Value

If `summary.fun = NULL` the function returns a list whose elements are the result of calling `spwb`. Otherwise, the function applies `summary.fun` to each simulation result and returns these summaries (actually, a call to `sapply` is done).

Author(s)

Miquel De Cáceres Ainsa, CTFC

See Also

[spwb](#), [summary.spwb](#)

Examples

```
## Not run:
#Load example data and species parameters
data(examplemeteo)
data(exampleforestMED)
data(SpParamsMED)

#Initialize input
examplesoil = soil(defaultSoilParams(2))
control = defaultControl()
x = forest2spwbInput(exampleforestMED,examplesoil, SpParamsMED, control)

#Perform sensitivity analysis
res = spwb_sensitivity(x, examplesoil, examplemeteo, latitude = 41, elevation = 100)

## End(Not run)
```

spwb_stress

Drought stress indices

Description

Allows calculating annual-based or monthly-based drought stress indices from `spwb` objects.

Usage

```
spwb_stress(x, index = "NDD", freq = "years", bySpecies = FALSE, draw = TRUE)
```

Arguments

`x` An object of class `spwb`.

`index` A string with the index to be calculated, either "DI", "NDD", "ADS", "MDS" or "WSI" (see details).

freq	Frequency of stress statistics (see cut.Date). Normally, either "years" or "months" for yearly-based or monthly-based indices.
bySpecies	Allows aggregating output by species.
draw	A boolean flag to indicate that a plot should be returned.

Details

The currently available drought stress indices are:

- "ADS": Average of daily drought stress values for the period considered.
- "MDS": Maximum daily drought stress during the period considered.
- "DI": Drought intensity, as defined in De Cáceres et al. (2015).
- "NDD": Number of drought days, as defined in De Cáceres et al. (2015).
- "WSI": Water stress integral, as defined in Myers (1988).

Value

A data frame with periods (e.g., years or months) in rows and plant cohorts (or species) in columns. Values are the calculated stress index. If `draw=TRUE` a ggplot is returned instead.

Author(s)

Miquel De Cáceres Ainsa, CTFC

References

De Cáceres M, Martínez-Vilalta J, Coll L, Llorens P, Casals P, Poyatos R, Pausas JG, Brotons L. (2015) Coupling a water balance model with forest inventory data to predict drought stress: the role of forest structural changes vs. climate changes. *Agricultural and Forest Meteorology* 213: 77-90 (doi:10.1016/j.agrformet.2015.06.012).

Myers BJ (1988) Water stress integral - a link between short-term stress and long-term growth. *Tree Physiology* 4: 315–323 (doi: 10.1093/treephys/4.4.315)

See Also

[spwb](#), [summary.spwb](#)

spwb_validation	<i>Validation of soil-plant water balance simulations</i>
-----------------	---

Description

Compares model predictions against observed values

Usage

```
spwb_validation(x, measuredData, type = "SWC", cohort = NULL, draw = TRUE,
               plotType = "dynamics")
```

Arguments

x	An object of class <code>spwb</code> or <code>pwb</code> .
measuredData	A data frame with observed/measured values. Dates should be in row names, whereas columns should be named according to the type of output to be evaluated (see details).
type	A string with the kind of model output to be evaluated. Accepted values are "SWC" (soil moisture content), "REW" relative extractable water, "ETR" (total evapotranspiration), "E" (transpiration per leaf area) and "WP" (plant water potentials).
cohort	A string of the cohort to be compared (e.g. "T1_68"). If NULL results for the first cohort will be evaluated.
draw	A boolean flag to indicate that plots should be produced.
plotType	Plot type to draw, either "dynamics" or "scatter".

Details

Users should provide the appropriate columns in `measuredData`, depending on the type of output to be evaluated:

- "SWC" or "REW": A column named "SWC" should be present, containing soil moisture content in percent volume.
- "ETR": A column named "ETR" should be present, containing stand's evapotranspiration in mm/day. Observed values will be compared against modelled evapotranspiration (i.e. sum of transpiration, soil evaporation and interception loss) as well as against the sum of transpiration and soil evaporation only.
- "E": For each plant cohort whose transpiration is to be evaluated, a column starting with "E_" and continuing with a cohort name (e.g. "E_T1_68") with transpiration in mm/day on a leaf area basis.
- "WP": For each plant cohort whose transpiration is to be evaluated, two columns, one starting with "PD_" (for pre-dawn) and the other with "MD_" (for midday), and continuing with a cohort name (e.g. "PD_T1_68"). They should contain leaf water potential values in MPa. These are compared against sunlit water potentials.

Additional columns may exist with the standard error of measured quantities. These should be named as the referred quantity, followed by "_err" (e.g. "PD_T1_68_err"), and are used to draw confidence intervals around observations.

Value

A ggplot or evaluation statistics (a vector or a data frame depending on type):

- MAE: Mean absolute error.
- Bias: Mean deviation (positive values correspond to model overestimations).
- r: Pearson's linear correlation coefficient.
- NSE: Nash-Sutcliffe model efficiency coefficient.
- NSEabs: Modified Nash-Sutcliffe model efficiency coefficient (L1 norm) (Legates & McCabe 1999).

Author(s)

Miquel De Cáceres Ainsa, CTFC

References

Legates, D.R., McCabe, G.J., 1999. Evaluating the use of "goodness-of-fit" measures in hydrologic and hydroclimatic model validation. *Water Resour. Res.* 35, 233–241.

See Also

[spwb](#)

spwb_waterUseEfficiency

Water use efficiency

Description

Calculates water use efficiency (WUE), at different temporal scales, from soil plant water balance calculations

Usage

```
spwb_waterUseEfficiency(x, type = "Plant An/E", leaves = "average", freq = "days",  
draw = TRUE, ylim = NULL)
```


Arguments

x	An object of class spwb or pwb.
type	A string to indicate the scale of WUE calculation. Either: <ul style="list-style-type: none"> • "Leaf iWUE": Leaf intrinsic WUE, i.e. instantaneous ratio between photosynthesis and stomatal conductance (only for simulations with transpirationMode = "Sperry" and subdailyResults = TRUE). • "Leaf Ci": Leaf intercellular CO2 concentration (only for simulations with transpirationMode = "Sperry" and subdailyResults = TRUE). • "Plant An/E": Plant (cohort) photosynthesis over plant transpiration) • "Stand An/E": Stand photosynthesis over stand transpiration)
leaves	Either "sunlit", "shade" or "average". Refers to the WUE of different leaf types or the average (with weights according to the LAI of sunlit and shade leaves). Only relevant for type = "iWUE".
freq	Frequency of summary statistics (see cut.Date).
draw	A boolean flag to indicate that a plot should be returned.
ylim	Range of values for y.

Details

Temporal aggregation of WUE values is done differently depending on the value of type. For type = "Plant An/E" and type = "Stand An/E" sums or daily photosynthesis and transpiration are first calculated at the desired temporal scale and the ratio is calculated afterwards. For type = "Leaf iWUE" intrinsic WUE values are first calculated at the daily scale (as averages of instantaneous An/g_s ratios weighted by An) and then they are aggregated to the desired scale by calculating weighted averages, where weights are given by daily photosynthesis.

Value

If draw=TRUE a plot is returned. Otherwise, the function returns a matrix with WUE values, where rows are dates (at the desired temporal scale), and columns are plant cohorts. In the case of type = "Plant An/E" and type = "Stand An/E" values are in gC/L. In the case of type = "Leaf iWUE" values are in micromol of carbon per mmol of water.

Author(s)

Miquel De Cáceres Ainsa, CTFC

See Also

[spwb](#)

Stand values

Stand values

Description

Functions to calculate stand attributes of a [forest](#) object.

Usage

```
stand_basalArea(x)
stand_foliarBiomass(x, SpParams, gdd = NA, mode = "MED")
stand_fuel(x, SpParams, gdd = NA, includeDead = TRUE, mode = "MED")
stand_LAI(x, SpParams, gdd = NA, mode = "MED")
stand_phytovolume(x, SpParams)
```

Arguments

x	An object of class forest .
SpParams	A data frame with species parameters (see SpParamsMED).
gdd	Growth degree days (to account for leaf phenology effects).
includeDead	A flag to indicate that standing dead fuels (dead branches) are included.
mode	Calculation mode, either "MED" or "US".

Value

- `stand_basalArea`: Stand basal area (m²/ha).
- `stand_foliarBiomass`: Standing biomass of leaves (in kg/m²).
- `stand_fuel`: Stand fine fuel load (in kg/m²).
- `stand_LAI`: Stand leaf area index (m²/m²).
- `stand_phytovolume`: Stand shrub phytovolume (m³/m²).

Author(s)

Miquel De Cáceres Ainsa, CTFC

See Also

[forest](#), [plant_basalArea](#), [summary.forest](#)

Examples

```
#Default species parameterization
data(SpParamsMED)

#Load example plot
data(exampleforestMED)

#A short way to obtain total basal area
stand_basalArea(exampleforestMED)
```

supplyfunctions	<i>Hydraulic supply functions</i>
-----------------	-----------------------------------

Description

Set of functions used in the implementation of hydraulic supply functions (Sperry & Love 2015).

Usage

```
hydraulics_EXylem(psiPlant, psiUpstream,
                  kxylemmax, c, d, allowNegativeFlux = TRUE,
                  psiCav = 0)
hydraulics_EVanGenuchten(psiRhizo, psiSoil, krhizomax,
                          n, alpha, l = 0.5)
hydraulics_ECapacitance(psi, psiPrev, PLCprev,
                        V, fapo, c, d,
                        pi0, eps,
                        timestep)
hydraulics_ECrit(psiUpstream, kxylemmax, c, d, pCrit = 0.001)
hydraulics_E2psiXylem(E, psiUpstream,
                     kxylemmax, c, d, psiCav = 0)
hydraulics_E2psiXylemUp(E, psiDownstream,
                        kxylemmax, c, d, psiCav = 0)
hydraulics_E2psiVanGenuchten(E, psiSoil, krhizomax, n, alpha,
                              psiStep = -0.0001, psiMax = -10.0)
hydraulics_E2psiTwoElements(E, psiSoil, krhizomax, kxylemmax, n, alpha, c, d,
                             psiCav = 0, psiStep = -1e-04, psiMax = -10.0)
hydraulics_E2psiFineRootLeaf(E, psiFineRoot,
                              krootmax, rootc, rootd,
                              kstemmax, stemc, stemd,
                              kleafmax, leafc, leafd,
                              PLCstem)
hydraulics_E2psiBelowground(E, psiSoil,
                             krhizomax, nsoil, alphasoil,
                             krootmax, rootc, rootd,
                             psiIni = as.numeric(c(0)),
                             ntrial = 10, psiTol = 0.0001, ETol = 0.0001)
```



```

        PLCstem,
        minFlow = 0.0, maxNsteps=400,
        ETol = 0.0001, pCrit = 0.001)
hydraulics_supplyFunctionNetworkStem1(psiSoil,
        krhizomax, nsoil, alphasoil,
        krootmax, rootc, rootd,
        kstemmax, stemc, stemd,
        PLCstem,
        minFlow = 0.0, maxNsteps=400,
        ntrial = 200, psiTol = 0.0001, ETol = 0.0001, pCrit = 0.001)
hydraulics_supplyFunctionNetwork(psiSoil,
        krhizomax, nsoil, alphasoil,
        krootmax, rootc, rootd,
        kstemmax, stemc, stemd,
        kleafmax, leafc, leafd,
        PLCstem, minFlow = 0.0, maxNsteps=400,
        ntrial = 200, psiTol = 0.0001, ETol = 0.0001, pCrit = 0.001)

hydraulics_supplyFunctionPlot(x, soil, draw = TRUE, type = "E",
        speciesNames = FALSE, ylim=NULL)
hydraulics_regulatedPsiXylem(E, psiUpstream, kxylemmax, c, d, psiStep = -0.01)
hydraulics_regulatedPsiTwoElements(Emax, psiSoil, krhizomax, kxylemmax, n, alpha,
        c, d, dE = 0.1, psiMax = -10.0)

```

Arguments

v	Proportion of fine roots within each soil layer.
krhizomax	Maximum rhizosphere hydraulic conductance (defined as flow per leaf surface unit and per pressure drop).
kxylemmax	Maximum xylem hydraulic conductance (defined as flow per leaf surface unit and per pressure drop).
kleafmax	Maximum leaf hydraulic conductance (defined as flow per leaf surface unit and per pressure drop).
kstemmax	Maximum stem xylem hydraulic conductance (defined as flow per leaf surface unit and per pressure drop).
krootmax	Maximum root xylem hydraulic conductance (defined as flow per leaf surface unit and per pressure drop).
E	Flow per surface unit.
Emax	Maximum flow per surface unit.
Erootcrown	Flow per surface unit at the root crown.
psi	Water potential (in MPa).
psiPrev	Water potential (in MPa) in the previous time step.
psiDownstream	Water potential upstream (in MPa).
psiUpstream	Water potential upstream (in MPa). In a one-component model corresponds to soil potential. In a two-component model corresponds to the potential inside the roots.

psiCav	Minimum water potential (in MPa) experienced (for irreversible cavitation).
minFlow	Minimum flow in supply function.
psiPlant	Plant water potential (in MPa).
psiFineRoot	Water potential (in MPa) inside fine roots.
psiSoil	Soil water potential (in MPa). A scalar or a vector depending on the function.
psiRhizo	Soil water potential (in MPa) in the rhizosphere (root surface).
psiRootCrown	Soil water potential (in MPa) at the root crown.
psiStep	Water potential precision (in MPa).
psiTol	Precision for water potential estimates (in MPa).
psiIni	Vector of initial water potential values (in MPa).
psiMax	Minimum (maximum in absolute value) water potential to be considered (in MPa).
pCrit	Critical water potential (in MPa).
PLCstem	Proportion of loss conductance in the stem [0-1].
PLCprev	Previous proportion of loss conductance [0-1].
V	Capacity of the compartment per leaf area (in L/m ²).
fapo	Apoplastic fraction (proportion) in the segment.
pi0	Full turgor osmotic potential (MPa).
eps	Bulk modulus of elasticity (MPa).
dE	Increment of flow per surface unit.
ETol	Precision for water flow per surface unit.
c, d	Parameters of the Weibull function (generic xylem vulnerability curve).
rootc, rootd	Parameters of the Weibull function for roots (root xylem vulnerability curve).
stemc, stemd	Parameters of the Weibull function for stems (stem xylem vulnerability curve).
leafc, leafd	Parameters of the Weibull function for leaves (leaf vulnerability curve).
n, alpha, l	Parameters of the Van Genuchten function (rhizosphere vulnerability curve).
nsoil, alphasoil	Parameter vectors of the Van Genuchten function (rhizosphere vulnerability curve) with one value for each soil layer.
allowNegativeFlux	A boolean to indicate whether negative flux (i.e. from plant to soil) is allowed.
maxNsteps	Maximum number of steps in the construction of supply functions.
ntrial	Maximum number of steps in Newton-Raphson optimization.
x	An object of class <code>spwbInput</code> .
soil	A list containing the description of the soil (see <code>soil</code>).
type	Plot type for <code>hydraulics_supplyFunctionPlot</code> , either "E", "ERhizo", "psiStem", "psiRoot", "psiRhizo" or "dEdP".
draw	A flag to indicate whether the supply function should be drawn or just returned.
speciesNames	A flag to indicate the use of species names instead of cohort names in plots.
ylim	Graphical parameter to override function defaults.
timestep	Time step in seconds.

Details

Function `hydraulics_supplyFunctionPlot` draws a plot of the supply function for the given soil object and network properties of each plant cohort in `x`. Function `hydraulics_vulnerabilityCurvePlot` draws a plot of the vulnerability curves for the given soil object and network properties of each plant cohort in `x`.

Value

Values returned for each function are:

- `hydraulics_E2psiXylem`: The plant (leaf) water potential (in MPa) corresponding to the input flow, according to the xylem supply function and given an upstream (soil or root) water potential.
- `hydraulics_E2psiVanGenuchten`: The root water potential (in MPa) corresponding to the input flow, according to the rhizosphere supply function and given a soil water potential.
- `hydraulics_E2psiTwoElements`: The plant (leaf) water potential (in MPa) corresponding to the input flow, according to the rhizosphere and plant supply functions and given an input soil water potential.
- `hydraulics_E2psiNetwork`: The rhizosphere, root crown and plant (leaf) water potential (in MPa) corresponding to the input flow, according to the vulnerability curves of rhizosphere, root and stem elements in a network.
- `hydraulics_Ecrit`: The critical flow according to the xylem supply function and given an input soil water potential.
- `hydraulics_EVanGenuchten`: The flow (integral of the vulnerability curve) according to the rhizosphere supply function and given an input drop in water potential (soil and rhizosphere).
- `hydraulics_EXylem`: The flow (integral of the vulnerability curve) according to the xylem supply function and given an input drop in water potential (rhizosphere and plant).
- `hydraulics_supplyFunctionOneXylem`, `hydraulics_supplyFunctionTwoElements` and `hydraulics_supplyFunctionNetwork`: A list with different numeric vectors with information of the two-element supply function:
 - `E`: Flow values (supply values).
 - `FittedE`: Fitted flow values (for `hydraulics_supplyFunctionTwoElements`).
 - `Elayers`: Flow values across the roots of each soil layer (only for `hydraulics_supplyFunctionNetwork`).
 - `PsiRhizo`: Water potential values at the root surface (only for `hydraulics_supplyFunctionNetwork`).
 - `PsiRoot`: Water potential values inside the root crown (not for `hydraulics_supplyFunctionOneXylem`).
 - `PsiPlant`: Water potential values at the canopy (leaf).
 - `dEdP`: Derivatives of the supply function.
- `hydraulics_supplyFunctionPlot`: If `draw = FALSE` a list with the result of calling `hydraulics_supplyFunctionNetwork` for each cohort.
- `hydraulics_regulatedPsiXylem`: Plant water potential after regulation (one-element loss function) given an input water potential.
- `hydraulics_regulatedPsiTwoElements`: Plant water potential after regulation (two-element loss function) given an input soil water potential.

Author(s)

Miquel De Cáceres Ainsa, CTFC.

Arguments

psiSym, psiApo	Symplastic or apoplastic water potential (MPa).
RWC	Relative water content [0-1].
pi0	Full turgor osmotic potential (MPa).
epsilon	Bulk modulus of elasticity (MPa).
c,d	Parameters of the xylem vulnerability curve.
af	Apoplastic fraction (proportion) in the segment (e.g. leaf or stem).
femb	Fraction of embolized conduits.
x	An object of class <code>spwbInput</code> .
segment	Segment whose relative water content curve to plot, either "stem" or "leaf".
fraction	Tissue fraction, either "symplastic", "apoplastic" or "all".
psiVec	Vector of water potential values to evaluate for the pressure-volume curve.
speciesNames	A flag to indicate the use of species names instead of cohort names in plots.
density	Tissue density (g-cm ⁻¹).
d0	Matric density (g-cm ⁻¹).
spwb	Object returned by function <code>spwb</code> or <code>growth</code> .

Details

Details of the tissue moisture calculations are given in a vignette. Function `moisture_cohortFMC()` calculates the fuel moisture content of leaves and twigs of each cohort, from the results of soil water balance.

Value

Values returned for each function are:

- `moisture_symplasticRWC`: Relative water content [0-1] of the symplastic fraction.
- `moisture_apoplasticRWC`: Relative water content [0-1] of the apoplastic fraction.
- `moisture_symplasticWaterPotential`: Water potential (in MPa) of the symplastic fraction.
- `moisture_apoplasticWaterPotential`: Water potential (in MPa) of the apoplastic fraction.
- `moisture_turgorLossPoint`: Water potential (in MPa) corresponding to turgor loss point.
- `moisture_segmentRWC`: Segment relative water content [0-1].

Author(s)

Miquel De Cáceres Ainsa, CTFC

References

- Bartlett, M.K., Scoffoni, C., Sack, L. 2012. The determinants of leaf turgor loss point and prediction of drought tolerance of species and biomes: a global meta-analysis. *Ecology Letters* 15: 393–405.
- Hölttä, T., Cochard, H., Nikinmaa, E., Mencuccini, M. 2009. Capacitive effect of cavitation in xylem conduits: Results from a dynamic model. *Plant, Cell and Environment* 32: 10–21.
- Martin-StPaul, N., Delzon, S., Cochard, H. 2017. Plant resistance to drought depends on timely stomatal closure. *Ecology Letters* 20: 1437–1447.

See Also

[hydraulics_psi2K](#), [hydraulics_supplyFunctionPlot](#), [spwb](#), [soil](#)

Examples

```
psi = seq(-10,0, by=0.1)
rwc_s = rep(NA, length(psi))
for(i in 1:length(psi)) rwc_s[i] = moisture_symplasticRWC(psi[i],-3,12)
plot(psi, rwc_s, type="l", xlab="Water potential (MPa)", ylab = "Symplasmic RWC")
```

transpiration

Transpiration submodel functions

Description

Set of high-level functions used in the calculation of stomatal conductance and transpiration. Function `transp_profitMaximization` calculates gain and cost functions, as well as profit maximization from supply and photosynthesis input functions. Function `transp_stomatalRegulationPlot` produces a plot with the cohort supply functions against water potential and a plot with the cohort photosynthesis functions against water potential, both with the maximum profit values indicated.

Usage

```
transp_profitMaximization(supplyFunction, photosynthesisFunction, Gwmin, Gwmax,
  gainModifier = 1.0, costModifier = 1.0, costWater = "dEdP")
transp_transpirationGranier(x, soil, meteo, day,
  modifyInputX = TRUE, modifyInputSoil = TRUE)
transp_transpirationSperry(x, soil, meteo, day,
  latitude, elevation, slope, aspect,
  canopyEvaporation = 0, snowMelt = 0, soilEvaporation = 0,
  stepFunctions = NA,
  modifyInputX = TRUE, modifyInputSoil = TRUE)
transp_stomatalRegulationPlot(x, soil, meteo, day, timestep,
  latitude, elevation, slope = NA, aspect = NA,
  type="E")
```

Arguments

`supplyFunction` Water supply function (see [hydraulics_supplyFunctionNetwork](#)).

`photosynthesisFunction` Function returned by `photo_photosynthesisFunction()`.

`Gwmin`, `Gwmax` Minimum and maximum stomatal conductance (mol-m-2-s-1).

`gainModifier`, `costModifier` Modifiers (exponents) of the gain and cost functions defined in Sperry et al. (2017).

`costWater` Cost water function, either "dEdP" (default) or "kterm" (experimental).

x	An object of class <code>spwbInput</code> built using the 'Sperry' transpiration mode.
soil	An object of class <code>soil</code> .
meteo	A data frame with daily meteorological data series: <ul style="list-style-type: none"> • DOY: Day of the year (Julian day). • Precipitation: Precipitation (in mm). • MeanTemperature: Mean temperature (in degrees Celsius). • MinTemperature: Minimum temperature (in degrees Celsius). • MaxTemperature: Maximum temperature (in degrees Celsius). • MinRelativeHumidity: Minimum relative humidity (in percent). • MaxRelativeHumidity: Maximum relative humidity (in percent). • Radiation: Solar radiation (in MJ/m²/day). • WindSpeed: Wind speed (in m/s). If not available, this column can be left with NA values.
day	An integer to identify a day within meteo.
timestep	An integer between 1 and ndailysteps specified in x (see <code>defaultControl</code>).
latitude	Latitude (in degrees).
elevation, slope, aspect	Elevation above sea level (in m), slope (in degrees) and aspect (in degrees from North).
canopyEvaporation	Canopy evaporation (from interception) for day (mm).
soilEvaporation	Bare soil evaporation for day (mm).
snowMelt	Snow melt values for day (mm).
stepFunctions	An integer to indicate a simulation step for which photosynthesis and profit maximization functions are desired.
type	A string with plot type, either "E" (transpiration flow), "Ag" (gross photosynthesis), "An" (net photosynthesis), "Gw" (stomatal conductance), "T" (temperature) or "VPD" (vapour pressure deficit).
modifyInputX, modifyInputSoil	Boolean flags to indicate that the input x and soil objects are allowed to be modified during the simulation.

Value

Function `transp_transpirationGranier` and `transp_transpirationSperry` return a list with the following elements:

- "cohorts": A data frame with cohort information, copied from `spwbInput`.
- "Stand": A vector of stand-level variables.
- "Plants": A data frame of results for each plant cohort. When using `transp_transpirationGranier`, element "Plants" includes:
 - "LAI": Leaf area index of the plant cohort.

- "AbsorbedSWRFraction": Fraction of SWR absorbed by each cohort.
- "Transpiration": Transpired water (in mm) corresponding to each cohort.
- "Photosynthesis": Net photosynthesis (in gC/m²) corresponding to each cohort.
- "psi": Water potential (in MPa) of the plant cohort (average over soil layers).
- "DDS": Daily drought stress [0-1] (relative whole-plant conductance).

When using `transp_transpirationSperry`, element "Plants" includes:

- "LAI": Leaf area index of the plant cohort.
 - "Extraction": Water extracted from the soil (in mm) for each cohort.
 - "Transpiration": Transpired water (in mm) corresponding to each cohort.
 - "GrossPhotosynthesis": Gross photosynthesis (in gC/m²) corresponding to each cohort.
 - "NetPhotosynthesis": Net photosynthesis (in gC/m²) corresponding to each cohort.
 - "RootPsi": Minimum water potential (in MPa) at the root collar.
 - "StemPsi": Minimum water potential (in MPa) at the stem.
 - "StemPLC": Proportion of conductance loss in stem.
 - "LeafPsiMin": Minimum (predawn) water potential (in MPa) at the leaf (representing an average leaf).
 - "LeafPsiMax": Maximum (midday) water potential (in MPa) at the leaf (representing an average leaf).
 - "LeafPsiMin_SL": Minimum (predawn) water potential (in MPa) at sunlit leaves.
 - "LeafPsiMax_SL": Maximum (midday) water potential (in MPa) at sunlit leaves.
 - "LeafPsiMin_SH": Minimum (predawn) water potential (in MPa) at shade leaves.
 - "LeafPsiMax_SH": Maximum (midday) water potential (in MPa) at shade leaves.
 - "dEdP": Overall soil-plant conductance (derivative of the supply function).
 - "DDS": Daily drought stress [0-1] (relative whole-plant conductance).
 - "StemRWC": Relative water content of stem tissue (including symplasm and apoplasm).
 - "LeafRWC": Relative water content of leaf tissue (including symplasm and apoplasm).
 - "StemSympRWC": Relative water content of symplastic stem tissue.
 - "LeafSympRWC": Relative water content of symplastic leaf tissue.
 - "WaterBalance": Plant water balance (extraction - transpiration).
- "Extraction": A data frame with mm of water extracted from each soil layer (in columns) by each cohort (in rows).
The remaining items are only given by `transp_transpirationSperry`:
 - "EnergyBalance": When using the 'Sperry' transpiration mode, the model performs energy balance of the stand and 'EnergyBalance' is a list with the following:
 - "Temperature": A data frame with the temperature of the atmosphere ('Tatm'), canopy ('Tcan') and soil ('Tsoil.1', 'Tsoil.2', ...) for each time step.
 - "CanopyEnergyBalance": A data frame with the components of the canopy energy balance (in W/m²) for each time step.
 - "SoilEnergyBalance": A data frame with the components of the soil energy balance (in W/m²) for each time step.
 - "RhizoPsi": Minimum water potential (in MPa) inside roots, after crossing rhizosphere, per cohort and soil layer.

- "SunlitLeaves" and "ShadeLeaves": Data frames for sunlit leaves and shade leaves and the following columns per cohort:
 - "LAI": Cumulative leaf area index of sunlit/shade leaves.
 - "Vmax298": Average maximum carboxylation rate for sunlit/shade leaves.
 - "Jmax298": Average maximum electron transport rate for sunlit/shade leaves.
- "ExtractionInst": Water extracted by each plant cohort during each time step.
- "PlantsInst": A list with instantaneous (per time step) results for each plant cohort:
 - "E": A data frame with the cumulative transpiration (mm) for each plant cohort during each time step.
 - "Ag": A data frame with the cumulative gross photosynthesis (gC/m²) for each plant cohort during each time step.
 - "An": A data frame with the cumulative net photosynthesis (gC/m²) for each plant cohort during each time step.
 - "SunlitLeaves" and "ShadeLeaves": Lists with instantaneous (for each time step) results for sunlit leaves and shade leaves and the following items:
 - * "Abs_SWR": A data frame with instantaneous absorbed short-wave radiation (SWR).
 - * "Abs_LWR": A data frame with instantaneous absorbed long-wave radiation (LWR).
 - * "An": A data frame with instantaneous net photosynthesis (in micromol/m²/s).
 - * "Ci": A data frame with instantaneous intercellular CO₂ concentration (in ppm).
 - * "GW": A data frame with instantaneous stomatal conductance (in mmol/m²/s).
 - * "VPD": A data frame with instantaneous vapour pressure deficit (in kPa).
 - * "Temp": A data frame with leaf temperature (in degrees Celsius).
 - * "Psi": A data frame with leaf water potential (in MPa).
 - "dEdP": A data frame with the slope of the plant supply function (an estimation of whole-plant conductance).
 - "RootPsi": A data frame with root crown water potential (in MPa) for each plant cohort during each time step.
 - "StemPsi": A data frame with stem water potential (in MPa) for each plant cohort during each time step.
 - "LeafPsi": A data frame with leaf (average) water potential (in MPa) for each plant cohort during each time step.
 - "StemPLC": A data frame with the proportion loss of conductance [0-1] for each plant cohort during each time step.
 - "StemRWC": A data frame with the (average) relative water content of stem tissue [0-1] for each plant cohort during each time step.
 - "LeafRWC": A data frame with the relative water content of leaf tissue [0-1] for each plant cohort during each time step.
 - "StemSympRWC": A data frame with the (average) relative water content of symplastic stem tissue [0-1] for each plant cohort during each time step.
 - "LeafSympRWC": A data frame with the relative water content of symplastic leaf tissue [0-1] for each plant cohort during each time step.
 - "PWB": A data frame with plant water balance (extraction - transpiration).
- "LightExtinction": A list of information regarding radiation balance through the canopy, as returned by function [light_instantaneousLightExtinctionAbsortion](#).

- "WindExtinction": A numeric vector with the wind speed (in m/s) for each plant cohort.
- "SupplyFunctions": If stepFunctions is not missing, a list of supply functions, photosynthesis functions and profit maximization functions.

Function `transp_profitMaximization` returns a list with the following elements:

- Cost: Cost function [0-1].
- Gain: Gain function [0-1].
- Profit: Profit function [0-1].
- iMaxProfit: Index corresponding to maximum profit (starting from 0).

Author(s)

Miquel De Cáceres Ainsa, CTFC

References

Sperry, J. S., M. D. Venturas, W. R. L. Anderegg, M. Mencuccini, D. S. Mackay, Y. Wang, and D. M. Love. 2017. Predicting stomatal responses to the environment from the optimization of photosynthetic gain and hydraulic cost. *Plant Cell and Environment* 40, 816-830 (doi: 10.1111/pce.12852).

See Also

[hydraulics_supplyFunctionNetwork](#), [biophysics_leafTemperature](#), [photo_photosynthesis](#), [spwb_day](#), [plot.spwb_day](#)

Examples

```
#Load example daily meteorological data
data(examplemeteo)

#Load example plot plant data
data(exampleforestMED)

#Default species parameterization
data(SpParamsMED)

#Initialize soil with default soil params (2 layers)
examplesoil = soil(defaultSoilParams(2))

#Initialize control parameters
control = defaultControl()

#Initialize input
x1 = forest2spwbInput(exampleforestMED,examplesoil, SpParamsMED, control)

# Transpiration according to Granier's model, plant water potential
# and plant stress for a given day
t1 = transp_transpirationGranier(x1, examplesoil, examplemeteo, 1,
                                modifyInputX = FALSE, modifyInputSoil = FALSE)
```

```

#Switch to 'Sperry' transpiration mode
control$transpirationMode="Sperry"

#Initialize soil with default soil params (2 layers)
examplesoil2 = soil(defaultSoilParams(2))

#Initialize input
x2 = forest2spwbInput(exampleforestMED,examplesoil2, SpParamsMED, control)

# Transpiration according to Sperry's model
t2 = transp_transpirationSperry(x2, examplesoil2, examplometeo, 1,
                               latitude = 41.82592, elevation = 100, slope = 0, aspect = 0,
                               modifyInputX = FALSE, modifyInputSoil = FALSE)

# Stomatal VPD curve and chosen value for the 12th time step at day 100
transp_stomatalRegulationPlot(x2, examplesoil2, examplometeo, day=100, timestep = 12,
                              latitude = 41.82592, elevation = 100, type="VPD")

```

Vertical profiles *Vertical profiles*

Description

Functions to generate vertical profiles generated by an input [forest](#) object.

Usage

```

vprofile_leafAreaDensity(x, SpParams = NULL, z = NULL, gdd = NA, mode = "MED",
                        byCohorts = FALSE, bySpecies = FALSE, draw = TRUE,
                        xlim = NULL)
vprofile_rootDistribution(x, SpParams, d = NULL, bySpecies = FALSE, draw = TRUE,
                        xlim = NULL)
vprofile_fuelBulkDensity(x, SpParams, z = NULL, gdd = NA, mode = "MED",
                        draw = TRUE, xlim = NULL)
vprofile_PARExtinction(x, SpParams, z = NULL, gdd = NA, mode = "MED",
                       draw = TRUE, xlim = c(0,100))
vprofile_SWRExtinction(x, SpParams, z = NULL, gdd = NA, mode = "MED",
                       draw = TRUE, xlim = c(0,100))
vprofile_windExtinction(x, SpParams, wind20H, z = NULL, gdd = NA, mode = "MED",
                       draw = TRUE, xlim = NULL)

```

Arguments

x	An object of class forest
SpParams	A data frame with species parameters (see SpParamsMED).
z	A numeric vector with height values.
d	A numeric vector with soil layer widths.

<code>gdd</code>	Growth degree days.
<code>mode</code>	Calculation mode, either "MED" or "US".
<code>byCohorts</code>	Separate profiles for each cohort.
<code>bySpecies</code>	Aggregate cohort profiles by species.
<code>wind20H</code>	The value of measured wind speed at 6 m = 20ft (in m/s).
<code>draw</code>	Logical flag to indicate that a plot is desired.
<code>xlim</code>	Limits of the x-axis.

Value

A numeric vector with values measured at each height. Units depend on the profile function:

- `vprofile_leafAreaDensity`: Cumulative LAI (m²/m²) per height bin.
- `vprofile_fuelBulkDensity`: Fuel bulk density (kg/m³) per height bin.
- `vprofile_PARExtinction`: Percent of photosynthetically active radiation (%) corresponding to each height.
- `vprofile_SWRExtinction`: Percent of shortwave radiation (%) corresponding to each height.
- `vprofile_windExtinction`: Wind speed (m/s) corresponding to each height.

Author(s)

Miquel De Cáceres Ainsa, CTFC

See Also

[forest](#)

Examples

```
#Default species parameterization
data(SpParamsMED)

#Load example plot plant data
data(exampleforestMED)

vprofile_leafAreaDensity(exampleforestMED, SpParamsMED)
vprofile_fuelBulkDensity(exampleforestMED, SpParamsMED)

vprofile_PARExtinction(exampleforestMED, SpParamsMED)
vprofile_SWRExtinction(exampleforestMED, SpParamsMED)

vprofile_windExtinction(exampleforestMED, SpParamsMED, 20)
```


Index

* datasets

- exampleforest, [12](#)
- examplemeteo, [13](#)
- SFM_metric, [59](#)
- SpParams, [72](#)

- biophysics, [3](#)
- biophysics_leafTemperature, [44](#), [110](#)
- biophysics_leafTemperature (biophysics), [3](#)
- biophysics_radiationDiurnalPattern (biophysics), [3](#)
- biophysics_temperatureDiurnalPattern (biophysics), [3](#)
- biophysics_waterDynamicViscosity (biophysics), [3](#)

- carbon, [4](#)
- carbon_leafStarchCapacity (carbon), [4](#)
- carbon_leafStructuralBiomass (carbon), [4](#)
- carbon_osmoticWaterPotential (carbon), [4](#)
- carbon_relativeSapViscosity (carbon), [4](#)
- carbon_sapwoodStarchCapacity (carbon), [4](#)
- carbon_sapwoodStructuralBiomass (carbon), [4](#)
- carbon_sapwoodStructuralLivingBiomass (carbon), [4](#)
- carbon_sugarConcentration (carbon), [4](#)
- carbon_sugarStarchDynamicsLeaf (carbon), [4](#)
- carbon_sugarStarchDynamicsStem (carbon), [4](#)
- conductancefunctions, [6](#)
- cut.Date, [48](#), [94](#), [97](#)

- Date, [25](#), [76](#)
- defaultControl, [8](#), [12](#), [14](#), [26](#), [35](#), [78](#), [107](#)
- defaultSoilParams, [11](#), [38](#), [61](#), [62](#), [71](#)

- emptyforest (forest), [18](#)

- exampleforest, [12](#)
- exampleforestMED, [19](#)
- exampleforestMED (exampleforest), [12](#)
- exampleforestUS (exampleforest), [12](#)
- examplemeteo, [13](#)
- extractSubdaily, [14](#)

- fire_behaviour, [15](#)
- fire_FCCS, [24](#)
- fire_FCCS (fire_behaviour), [15](#)
- fire_Rothermel, [60](#)
- fire_Rothermel (fire_behaviour), [15](#)
- forest, [12](#), [18](#), [20–22](#), [32](#), [34](#), [35](#), [38](#), [44–46](#), [71](#), [72](#), [80](#), [98](#), [111](#), [112](#)
- Forest manipulation, [20](#)
- forest2aboveground (modelInput), [34](#)
- forest2belowground (modelInput), [34](#)
- forest2growthInput (modelInput), [34](#)
- forest2spwbInput, [12](#), [56](#)
- forest2spwbInput (modelInput), [34](#)
- forest_mapShrubTable (forest_mapWoodyTables), [21](#)
- forest_mapTreeTable (forest_mapWoodyTables), [21](#)
- forest_mapWoodyTables, [19](#), [21](#)
- forest_mergeTrees (Forest manipulation), [20](#)
- fuel_FCCS, [15](#), [17](#)
- fuel_FCCS (fuel_properties), [22](#)
- fuel_properties, [22](#)
- fuel_stratification (fuel_properties), [22](#)
- fuel_windAdjustmentFactor (fuel_properties), [22](#)

- growth, [5](#), [14](#), [24](#), [34](#), [40](#), [47](#), [49](#), [83](#), [105](#)
- growth_day, [26](#), [27](#), [50](#)
- growth_day (spwb_day), [81](#)
- growthInput, [25](#), [27](#), [39](#), [53](#), [54](#), [82](#), [83](#)
- growthInput (modelInput), [34](#)

- hydraulics_averagePsi
(conductancefunctions), 6
- hydraulics_averageRhizosphereResistancePercent
(scalingconductance), 57
- hydraulics_correctConductanceForViscosity
(conductancefunctions), 6
- hydraulics_E2psiAboveground
(supplyfunctions), 99
- hydraulics_E2psiBelowground
(supplyfunctions), 99
- hydraulics_E2psiFineRootLeaf
(supplyfunctions), 99
- hydraulics_E2psiNetwork
(supplyfunctions), 99
- hydraulics_E2psiNetworkStem1
(supplyfunctions), 99
- hydraulics_E2psiTwoElements
(supplyfunctions), 99
- hydraulics_E2psiVanGenuchten
(supplyfunctions), 99
- hydraulics_E2psiXylem
(supplyfunctions), 99
- hydraulics_E2psiXylemUp
(supplyfunctions), 99
- hydraulics_ECapacitance
(supplyfunctions), 99
- hydraulics_ECrit (supplyfunctions), 99
- hydraulics_EVanGenuchten
(supplyfunctions), 99
- hydraulics_EXylem (supplyfunctions), 99
- hydraulics_findRhizosphereMaximumConductance
(scalingconductance), 57
- hydraulics_K2Psi
(conductancefunctions), 6
- hydraulics_leafWaterCapacity
(scalingconductance), 57
- hydraulics_maximumSoilPlantConductance
(scalingconductance), 57
- hydraulics_maximumStemHydraulicConductance,
8, 104
- hydraulics_maximumStemHydraulicConductance
(scalingconductance), 57
- hydraulics_psi2K, 59, 104, 106
- hydraulics_psi2K
(conductancefunctions), 6
- hydraulics_psi2Weibull
(conductancefunctions), 6
- hydraulics_psiCrit
(conductancefunctions), 6
- hydraulics_referenceConductivityHeightFactor
(scalingconductance), 57
- hydraulics_regulatedPsiTwoElements
(supplyfunctions), 99
- hydraulics_regulatedPsiXylem
(supplyfunctions), 99
- hydraulics_rootxylemConductanceProportions
(scalingconductance), 57
- hydraulics_soilPlantResistances, 91
- hydraulics_soilPlantResistances
(scalingconductance), 57
- hydraulics_stemWaterCapacity
(scalingconductance), 57
- hydraulics_supplyFunctionAboveground
(supplyfunctions), 99
- hydraulics_supplyFunctionBelowground
(supplyfunctions), 99
- hydraulics_supplyFunctionFineRootLeaf
(supplyfunctions), 99
- hydraulics_supplyFunctionNetwork, 44,
106, 110
- hydraulics_supplyFunctionNetwork
(supplyfunctions), 99
- hydraulics_supplyFunctionNetworkStem1
(supplyfunctions), 99
- hydraulics_supplyFunctionOneXylem
(supplyfunctions), 99
- hydraulics_supplyFunctionPlot, 8, 59,
106
- hydraulics_supplyFunctionPlot
(supplyfunctions), 99
- hydraulics_supplyFunctionThreeElements
(supplyfunctions), 99
- hydraulics_supplyFunctionTwoElements
(supplyfunctions), 99
- hydraulics_taperFactorSavage
(scalingconductance), 57
- hydraulics_terminalConduitRadius
(scalingconductance), 57
- hydraulics_vanGenuchtenConductance
(conductancefunctions), 6
- hydraulics_vulnerabilityCurvePlot
(conductancefunctions), 6
- hydraulics_xylemConductance
(conductancefunctions), 6
- hydraulics_xylemPsi
(conductancefunctions), 6

- hydrology_erFactor
(hydrology_rainInterception),
28
- hydrology_infiltrationAmount (soil
hydrology), 63
- hydrology_infiltrationRepartition
(soil hydrology), 63
- hydrology_interceptionPlot
(hydrology_rainInterception),
28
- hydrology_rainInterception, 28, 31
- hydrology_snowMelt (soil hydrology), 63
- hydrology_soilEvaporation, 31
- hydrology_soilEvaporation (soil
hydrology), 63
- hydrology_soilEvaporationAmount (soil
hydrology), 63
- hydrology_soilInfiltrationPercolation
(hydrology_soilWaterInputs), 29
- hydrology_soilWaterInputs, 29, 64

- light, 31
- light_cohortAbsorbedSWRFraction
(light), 31
- light_cohortSunlitShadeAbsorbedRadiation
(light), 31
- light_instantaneousLightExtinctionAbsortion,
83, 109
- light_instantaneousLightExtinctionAbsortion
(light), 31
- light_layerIrradianceFraction (light),
31
- light_layerIrradianceFractionBottomUp
(light), 31
- light_layerSunlitFraction (light), 31

- modelInput, 34
- modifyCohortParams (modifySpParams), 39
- modifySpParams, 39
- moisture_apoplasticPsi
(tissuemoisture), 104
- moisture_apoplasticRWC
(tissuemoisture), 104
- moisture_cohortFMC (tissuemoisture), 104
- moisture_pressureVolumeCurvePlot
(tissuemoisture), 104
- moisture_symplasticPsi
(tissuemoisture), 104

- moisture_symplasticRWC
(tissuemoisture), 104
- moisture_tissueFMC (tissuemoisture), 104
- moisture_tissueRWC (tissuemoisture), 104
- moisture_turgorLossPoint
(tissuemoisture), 104

- pheno_leafDevelopmentStatus
(pheno_updateLeaves), 40
- pheno_leafSenescenceStatus
(pheno_updateLeaves), 40
- pheno_updateLeaves, 40
- pheno_updatePhenology
(pheno_updateLeaves), 40
- photo, 41
- photo_electronLimitedPhotosynthesis
(photo), 41
- photo_GammaTemp (photo), 41
- photo_JmaxTemp (photo), 41
- photo_KmTemp (photo), 41
- photo_leafPhotosynthesisFunction
(photo), 41
- photo_multilayerPhotosynthesisFunction
(photo), 41
- photo_photosynthesis, 110
- photo_photosynthesis (photo), 41
- photo_rubiscolimitedPhotosynthesis
(photo), 41
- photo_sunshadePhotosynthesisFunction
(photo), 41
- photo_VmaxTemp (photo), 41
- Plant values, 44
- plant_basalArea, 72, 98
- plant_basalArea (Plant values), 44
- plant_characterParameter (Plant
values), 44
- plant_cover (Plant values), 44
- plant_crownBaseHeight (Plant values), 44
- plant_crownLength (Plant values), 44
- plant_crownRatio (Plant values), 44
- plant_density, 35
- plant_density (Plant values), 44
- plant_equilibriumLeafLitter (Plant
values), 44
- plant_equilibriumSmallBranchLitter
(Plant values), 44
- plant_foliarBiomass (Plant values), 44
- plant_fuel (Plant values), 44
- plant_height (Plant values), 44

- plant_ID, [35](#), [38](#)
- plant_ID (Plant values), [44](#)
- plant_LAI (Plant values), [44](#)
- plant_largerTreeBasalArea (Plant values), [44](#)
- plant_parameter (Plant values), [44](#)
- plant_phytovolume (Plant values), [44](#)
- plant_shrubIndividualArea (Plant values), [44](#)
- plant_species (Plant values), [44](#)
- plant_speciesName (Plant values), [44](#)
- plot.growth, [27](#)
- plot.growth (plot.spwb), [46](#)
- plot.growth_day, [83](#)
- plot.growth_day (plot.spwb_day), [50](#)
- plot.pwb (plot.spwb), [46](#)
- plot.pwb_day (plot.spwb_day), [50](#)
- plot.spwb, [46](#), [52](#), [80](#), [91](#)
- plot.spwb_day, [47](#), [50](#), [83](#), [110](#)
- print, [19](#)
- print.soil (soil), [61](#)
- print.summary.forest (forest), [18](#)
- pwb, [14](#), [34](#), [47](#), [48](#), [95](#)
- pwb (spwb), [76](#)

- redefineSoilLayers (soil), [61](#)
- resetInputs, [38](#), [53](#)
- root, [54](#)
- root_coarseRootLengths (root), [54](#)
- root_coarseRootLengthsFromVolume (root), [54](#)
- root_coarseRootSoilVolume (root), [54](#)
- root_coarseRootSoilVolumeFromConductance (root), [54](#)
- root_conicDistribution (root), [54](#)
- root_fineRootAreaIndex (root), [54](#)
- root_fineRootBiomass (root), [54](#)
- root_fineRootHalfDistance (root), [54](#)
- root_fineRootRadius (root), [54](#)
- root_fineRootSoilVolume (root), [54](#)
- root_horizontalProportions (root), [54](#)
- root_individualRootedGroundArea (root), [54](#)
- root_ldrDistribution, [85](#), [86](#), [88](#)
- root_ldrDistribution (root), [54](#)
- root_rhizosphereMaximumConductance (root), [54](#)
- root_specificRootSurfaceArea (root), [54](#)

- sapply, [92](#), [93](#)
- scalingconductance, [57](#)
- SFM_metric, [59](#)
- soil, [6](#), [8](#), [11](#), [12](#), [25](#), [30](#), [35](#), [38](#), [53](#), [56](#), [59](#), [61](#), [63](#), [66](#), [68–71](#), [76](#), [77](#), [82](#), [85–88](#), [90](#), [92](#), [102](#), [104](#), [106](#), [107](#)
- soil hydrology, [63](#)
- soil texture and hydraulics, [65](#)
- soil thermodynamics, [68](#)
- soil_psi (soil texture and hydraulics), [65](#)
- soil_psi2thetaSX, [62](#)
- soil_psi2thetaSX (soil texture and hydraulics), [65](#)
- soil_psi2thetaVG, [62](#)
- soil_psi2thetaVG (soil texture and hydraulics), [65](#)
- soil_retentionCurvePlot (soil texture and hydraulics), [65](#)
- soil_rockWeight2Volume (soil texture and hydraulics), [65](#)
- soil_saturatedConductivitySX, [62](#)
- soil_saturatedConductivitySX (soil texture and hydraulics), [65](#)
- soil_temperatureChange (soil thermodynamics), [68](#)
- soil_temperatureGradient (soil thermodynamics), [68](#)
- soil_thermalCapacity (soil thermodynamics), [68](#)
- soil_thermalConductivity (soil thermodynamics), [68](#)
- soil_theta (soil texture and hydraulics), [65](#)
- soil_theta2psiSX (soil texture and hydraulics), [65](#)
- soil_theta2psiVG (soil texture and hydraulics), [65](#)
- soil_thetaFC (soil texture and hydraulics), [65](#)
- soil_thetaSAT (soil texture and hydraulics), [65](#)
- soil_thetaSATSX (soil texture and hydraulics), [65](#)
- soil_thetaWP (soil texture and hydraulics), [65](#)
- soil_USDAType (soil texture and hydraulics), [65](#)

- soil_vanGenuchtenParamsCarsel (soil texture and hydraulics), 65
- soil_vanGenuchtenParamsToth (soil texture and hydraulics), 65
- soil_water (soil texture and hydraulics), 65
- soil_waterExtractable (soil texture and hydraulics), 65
- soil_waterFC (soil texture and hydraulics), 65
- soil_waterSAT (soil texture and hydraulics), 65
- soil_waterTableDepth (soil texture and hydraulics), 65
- soil_waterWP (soil texture and hydraulics), 65
- soilgridsParams, 12, 70
- Species values, 71
- species_basalArea (Species values), 71
- species_cover (Species values), 71
- species_density (Species values), 71
- species_foliarBiomass (Species values), 71
- species_fuel (Species values), 71
- species_LAI (Species values), 71
- species_phytovolume (Species values), 71
- SpParams, 72
- SpParamsMED, 12, 19, 21, 22, 28, 32, 35, 38–40, 45, 71, 98, 111
- SpParamsMED (SpParams), 72
- SpParamsUS (SpParams), 72
- spwb, 4, 8, 9, 11–14, 20, 24–26, 29, 33, 34, 38, 41, 44, 46–48, 50, 52, 54, 56, 59, 62, 64, 72, 75, 76, 82–84, 86–88, 90–97, 104–106
- spwb_day, 9, 31, 50, 52, 78, 80, 81, 110
- spwb_ldrCalibration, 84
- spwb_ldrExploration (spwb_ldrOptimization), 86
- spwb_ldrOptimization, 56, 80, 86, 86
- spwb_maximumTranspirationRatioPlot, 89
- spwb_resistances, 91
- spwb_sensitivity, 92
- spwb_stress, 93
- spwb_validation, 95
- spwb_waterUseEfficiency, 96
- spwbInput, 6, 9, 11, 28, 39–41, 53, 54, 76, 77, 80, 82, 83, 85, 87, 90, 92, 102, 105, 107
- spwbInput (modelInput), 34
- Stand values, 98
- stand_basalArea (Stand values), 98
- stand_foliarBiomass (Stand values), 98
- stand_fuel (Stand values), 98
- stand_LAI (Stand values), 98
- stand_phytovolume (Stand values), 98
- summary, 19
- summary.forest, 20, 46, 72, 98
- summary.forest (forest), 18
- summary.growth (plot.spwb), 46
- summary.pwb (plot.spwb), 46
- summary.spwb, 93, 94
- summary.spwb (plot.spwb), 46
- supplyfunctions, 99
- tissuemoisture, 104
- transp_profitMaximization (transpiration), 106
- transp_stomatalRegulationPlot (transpiration), 106
- transp_transpirationGranier, 77, 82
- transp_transpirationGranier (transpiration), 106
- transp_transpirationSperry, 50, 77, 82, 83
- transp_transpirationSperry (transpiration), 106
- transpiration, 106
- Vertical profiles, 111
- vprofile_fuelBulkDensity (Vertical profiles), 111
- vprofile_leafAreaDensity (Vertical profiles), 111
- vprofile_PARExtinction (Vertical profiles), 111
- vprofile_rootDistribution (Vertical profiles), 111
- vprofile_SWRExtinction (Vertical profiles), 111
- vprofile_windExtinction (Vertical profiles), 111